

Plant Disease Reporter

Crops Research Division
Agricultural Research Service

UNITED STATES DEPARTMENT OF AGRICULTURE

VOLUME 50

NUMBER 13

1966 Index
(Issued April 1967)



The Plant Disease Reporter is published monthly as a service to American agriculture. The articles emphasize new developments in plant pathology, including disease incidence and severity, serious outbreaks and epidemics, investigative techniques, and control chemicals. The rapid issuance of such information permits plant pathologists to recommend the judicious use of pesticides or other appropriate control measures against impending disease epidemics. The Crops Research Division, in publishing this material, does not assume responsibility for the subject matter.

Issued by

CROPS RESEARCH DIVISION

Plant Industry Station
Volume 50

Beltsville, Maryland
Number 13

AUTHOR INDEX

- | | |
|--|-------------------------------|
| ADAMS, R. E., (545) | BRAVO, LUIS E., (101) |
| ADLERZ, W. C., (602) | BRETZ, T. W., (45) |
| AFANASIEV, M. M., 928 | BRINKERHOFF, L. A., 323 |
| AGNIHOTRI, J. P., 921 | BROWDER, L. E., 673, (756) |
| AGRIOS, GEORGE N., 151 | BROWN, G. ELDON, 860 |
| AICHELE, MURIT D., (3) | BRZESKI, M. W., (584) |
| ALLAM, E. K., 761 | BUCHHOLTZ, W. F., 88 |
| ALLAN, R. E., (205) | BUGBEE, W. M., 576 |
| ALLEN, ROSS M., 145 | BUTLER, E. E., 796 |
| ALLEN, W. WAYNE, 622 | BYRD, D. W., Jr., 954 |
| ANDERSON, P. M., (363) | BYRNE, T. G., (751) |
| ARMSTRONG, G. M., 391 | |
| ARMSTRONG, JOANNE K., (391) | CALVERT, OSCAR H., (753) |
| ARNY, D. C., (12), (377), 380 | CAMPBELL, W. A., (393) |
| ATLEE, C. B., Jr., (611) | CANNON, T. F., (664) |
| AULAKH, KIRPAL S., 314 | CARLSON, L. W., 539 |
| | CARPENTER, JOHN B., (145) |
| BACHELDER, S., (294) | CASTIBLANCO, LUIS E., (101) |
| BAGGETT, J. R., 532 | CEPONIS, MICHAEL J., 222 |
| BAIN, DOUGLAS C., 770, (770) | CHAKRAVARTI, B. P., 701 |
| BAJAJ, B. S., (299) | CHAMBERS, ALBERT Y., (608) |
| BALDRIDGE, JOE D., (753) | CHI, C. C., 451, 695 |
| BALL, ELLEN M., (951) | CHILDERS, W. R., (695) |
| BANTTARI, E. E., 17 | CHILDS, J. F. L., 81 |
| BARBE, G. DOUGLAS, 768 | CLARK, W. A., (502) |
| BARD, G., (722) | COHEN, ELIAHOU, (251), (867) |
| BARKAI-GOLAN, RIVKA, 874 | COHEN, Y., (635) |
| BARKER, K. R., (420), (954), 957 | COHN, E., 711 |
| BARNES, D. K., (484) | COLE, H., Jr., 494 |
| BARNES, GEORGE L., 599 | COLLINS, DON D., 45 |
| BASHAW, E. C., (622) | COMBS, R. L., Jr., (570) |
| BATRA, S. K., 299 | CONVERSE, R. H., 351 |
| BAUER, A. H., (6) | COOK, A. A., (761) |
| BEAN, GEORGE A., 942 | COOPER, R. L., (557) |
| BEARDMORE, R. W., (798) | COOPER, W. E., 136 |
| BEGA, ROBERT V., (181), (245), 439,
832 | CORDELL, CHARLES E., 589 |
| BELL, DURHAM K., 162 | COUNTER, BEN F., (371) |
| BERAHA, L., (283) | COVEY, R. A., (645) |
| BERGER, RICHARD D., (701) | COVEY, RONALD P., Jr., 787 |
| BERKENKAMP, B., 390 | COX, R. S., 124, (354), 699 |
| BERRY, F. H., 45 | COYNE, D. P., 29 |
| BIESBROCK, J. A., 360 | CRANE, J. L., 464 |
| BIRCHFIELD, WRAY, 149, (500) | CRITTENDEN, H. W., (464), 910 |
| BLODGETT, EARLE C., 3 | CROSIER, WILLARD F., (321) |
| BOND, WILLIAM P., 325 | CROSS, J. E., 557, (560) |
| BOONE, DONALD M., (539), 543, (957) | CROSSAN, D. F., (84), (647) |
| BORDERS, HUEY I., 847 | CULP, T. W., 35 |
| BOYCE, JOHN S., Jr., (360), 482 | |
| BOYLE, LYTTON W., (810) | DAFT, GILBERT C., 493 |
| BRAKKE, MYRON K., (951) | DALE, J. L., 441 |
| BRAUN, ALVIN J., 783, (877) | DAVIS, D. R., 906 |
| BRAVERMAN, SAMUEL W., 321, 488,
849 | DAVIS, J. R., (53) |
| | DÍAZ, FRANCISCO, (422) |
| | DICKENS, LESTER E., (371) |

- DICKERSON, O. J., 396
DICKSON, R. C., 305
DIMOCK, A. W., 751
DISHON, IRENE, (21)
DOBIE, N. D., 426
DOEPPEL, R. F., 494
DOOLEY, H. L., (97)
DOOLING, O. J., 862
DOTTA, D., (439)
DOUGLAS, W. A., (570)
DUBIN, H. JESSE, 280
DUKES, P. D., (275), (444), (520), (836)
DUNLEAVY, JOHN, (598)
DWINELL, L. D., (406)
- ECK, PAUL, (346)
ECKESS, E. D., (571)
EDGERTON, L. J., (240)
EDMUNDS, J. E., 478
EL-HAIDARI, H. S., 960
ELLINGBOE, A. H., (190)
ENGEL, R. E., (703)
ENGLISH, HARLEY, (53)
EPPS, JAMES M., 608
EPSTEIN, A. H., 167, 420
ERWIN, D. C., (930)
EVERETT, PAUL H., (340)
- FARRELL, K. M., (478)
FATHI, SAAD M., (401)
FEDER, WILLIAM A., 429
FILER, T. H., Jr., 264, 525, 751
FISHER, FRAN. E., 357
FLEISCHMANN, G., 750
FLIEGEL, PETER, 240
FORD, RICHARD E., (88), (814)
FOSKETT, RICHARD L., (309)
FOSTER, R. E., (49), 79
FRAZIER, D. H., (510)
FRAZIER, N. W., 318
FRAZIER, W. A., (532)
FREDERIKSEN, R. A., 840
FREEMAN, T. E., 292
FREITAG, J. H., (318)
FREYRE, R. H., (484)
FRIDLUND, PAUL R., 740, 902
FROSHEISER, F. I., 435
FULTON, N. D., (350)
FULTON, ROBERT W., (613)
FUNK, C. R., (294), (703)
FUTRELL, MAURICE C., (373), 606,
641, 828
- GARREN, R. E., (843)
GARRETT, WILEY N., (839)
GASKILL, JOHN O., 457
GASKIN, T. A., 430
GAUTREAU, E. J., (419)
GHAFFAR, ABDUL, 171
GILMER, R. M., 461
- GOLDEN, A. MORGAN, (645), (798)
GOOCH, F. S., 525
GOODING, G. V., Jr., 308
GOTH, R. W., 33, 110
GRAVES, ALFRED A., Jr., (482)
GREATHEAD, A. S., 327
GRIERSON, W., (860)
GRIFFITHS, D. A., 116, 261
GROVES, A. B., (461)
GUDAUSKAS, ROBERT T., 885
GUENGERICH, H. W., 367
GUTTER, Y., 782
- HAASIS, F. A., (664)
HAGEDORN, D. J., (91), (578)
HALISKY, P. M., 294, 703
HALL, D. H., 793, (796)
HALLIWELL, ROBERT S., 75
HAMBLEN, M. L., (15)
HAMILTON, R. I., (822)
HAMMONS, R. O., (520)
HANLIN, RICHARD T., 306
HANSEN, A. JUERGEN, 229
HARDISON, JOHN R., 624
HARLAN, D. P., 548
HARMON, DANIEL L., (502)
HARMON, S. A., (443)
HARRISON, A. L., 855
HARRISON, MONTY D., 897
HARROW, K. M., (730)
HARTWIG, E. E., (104)
HARVEY, GEORGE M., 554
HASAN, S., 764
HASTINGS, K. M., (910)
HELTON, A. W., (773)
HENDRIX, FLOYD F., Jr., 393
HENDRIX, J. WALTER, (369)
HENSON, LAWRENCE, (218)
HERNANDEZ, TEME P., (500)
HERR, L. J., 777
HIBBEN, C. R., 905
HILTY, J. W., 427
HIMELICK, E. B., (473)
HINDS, THOMAS E., 741
HIRATSUKA, Y., 419
HOBBS, C. D., 373
HOCKING, D., 593
HOES, J. A., 62, 154
HOLDEMAN, Q. L., (318)
HOPKINS, DONALD L., (218)
HOPPE, PAUL E., 377
HORNE, C. WENDELL, 869
HORNOR, C. E., 97
HOWELL, R. W., (104)
HSI, DAVID C. H., 175, 352
HUBER, D. M., 852
HULLUKA, MENGISTU, 631
HUTCHINS, LEE M., 219
HUTCHINS, P. C., (429)
HYRE, R. A., 734

- IIDA, T. T., (453)
IZADPANAH, K., 779
- JEDLINSKI, H., (380)
JENKINS, L., (548)
JENKINS, S. F., Jr., (275), 520, 836
JENSEN, HAROLD J., 923
JENSEN, RAY E., 810
JOHNSON, D. E., (133), (611), 915
JOHNSON, E. M., 218
JOHNSON, LOWELL B., 108
JOHNSON, R. E., (81)
JOHNSTON, C. O., 756
JONES, JOHN PAUL, 340, 707
JORGENSEN, E. C., 650
JOSEPHSON, L. M., (427)
- KAHAN, R. S., (874)
KAIS, A. G., (388), 842
KALLIO, ARVO, 69
KALOOSTIAN, G. H., (270)
KANTZES, J. G., (351)
KAPPELMAN, A. J., Jr., 655
KATSANOS, R. A., 287
KATZIR, R., (594)
KEEN, NOEL T., (192)
KENNEDY, B. W., (557), 560
KEPLINGER, JOSEPH A., (783), (877)
KERNKAMP, M. F., (576), (800)
KHANNA, R. N., (127)
KILBY, W. W., (54), (717)
KILPATRICK, R. A., (622), 789
KLINE, D. M., (382)
KOENIGS, J. W., 528
KOMMEDAHL, THOR, 26
KOZELNICKY, GEORGE M., 839, 840
KRIKUN, J., (711)
KRULL, CHARLES F., 101
KUNTZ, JAMES E., (677), (736)
- LABORDE, J. A., (947)
LACKEY, C. F., 479
LAIRD, E. F., Jr., (305)
LAMBE, R. C., (88)
LAMBERT, J. W., (557)
LANGDON, K. R., (822)
LATHAM, A. J., 469
LAURENT, T. H., 921
LAUT, JOHN G., 208
LAWRENCE, J. J., (208)
LAYNE, R. E. C., 112
LEAR, BERT, 132, 133, 611, 858, (915)
LEATH, K. T., 312
LEBEN, CURT, (493)
LEEPER, PAUL W., (460)
LEWIS, MARJORIE H., 681
LEWIS, W. A., (54), (717)
LIM, W. C., (116), (261)
LITTRELL, R. H., 882
LIVINGSTON, CLARK H., (897)
LOEGERING, WILLIAM Q., 502
- LORIO, PETER L., Jr., 596
LUCHT, DONALD D., (126)
LUEPSCHEN, N. S., 565
LUKE, H. H., (292)
- MacKENZIE, D. R., (6), 363, 814
MacSWAN, I. C., (267)
MAIER, CHARLES R., 301
MANNING, W. J., 84, 647
MARLATT, ROBERT B., 687, 689
MARTENS, J. W., 12
MARTIN, JEAN E., 822
MARTIN, NEIL E., 369
MARTIN, W. J., 500
MATHRE, D. E., 930
MATHUR, S. B., 619
MAXIE, E. C., (327)
McCARTER, STATES M., (885)
McCARTNEY, WILLIAM O., (768)
McCLAIN, L. A., 112
McCLELLAN, W. D., 725
McCOLLOCH, LACY P., 178, 348
McCRUM, R. C., 40
McDANIEL, M. C., 350
McDONALD, H. H., (947)
McELROY, F. D., 581
McINTOSH, D. L., 267
McKINNEY, H. H., 951
McLEAN, D. M., 871
McWHORTER, F. P., (532)
MEYER, R. W., (938)
MICHEL, L. J., 935
MILBRATH, J. A., 59
MILLECAN, ARTHUR A., 211
MILLER, D. R., (439)
MILLER, J. W., 187
MILLER, LAWRENCE I., 139
MILLER, P. M., 201, 506
MILLER, P. W., (426), 843
MILLER, PAUL R., 803
MILLIKAN, D. F., (367)
MINK, G. I., (461)
MISHRA, M. D., (619)
MITCHELL, J. E., 91
MIYAGAWA, S. T., (132), (133), (611)
MOHAMED, HOSNI A., 401
MOLLER, W. J., 53
MONCRIEF, JAMES B., (840)
MONEIM, AHMED T. ABDEL, (401)
MOORE, D. M., (910)
MOORE, J. N., 105
MORGAN, FREDERIC L., 104, 598
MORRIS, C. L., 510
MORTON, D. J., 96, 275, (443), 444, (836)
MOTSINGER, R. E., 686
MUELLER, WALTER C., 899
MUKHERJEE, A. K., 88
- NATTI, J. J., 614
NEELY, DAN, 473

- NELSON, MERRITT R., (629), 947
 NELSON, R. R., 382
 NETZER, DAVID, 21, 594
 NEWHALL, ALLAN G., 422
 NICHOLS, LESTER P., 515
 NICKLE, W. R., 798
 NIELSEN, L. W., 730
 NITZANY, F. E., 158
 NORTON, DON C., 523
 NORTON, J. D., (661)
 NUSBAUM, C. J., (954)
- O'BANNON, JOHN H., (512)
 OHMAN, JOHN H., 402
 OLOFSSON, JAN, 257
 ORELLANA, R. G., 7
 ORLOB, GERT B., 819, 917
 ORR, CALVIN C., (149), 645
 OSHIMA, NAGAYOSHI, 309, 371
 OSTAZESKI, STANLEY A., 825
 OTTERBACHER, A. G., (65)
 OU, S. H., (453)
 OUELLETTE, G. B., 722
- PAI, H. SURENDRANATH, 313, 518
 PALMER, L. T., 234
 PALMITER, D. H., (783), 877
 PAPPELIS, A. J., (287)
 PARACER, S. M., 584
 PARKER, K. G., (240)
 PARRIS, G. K., 243
 PARTRIDGE, A. D., 497
 PATEL, K. P., (26)
 PATWARDHAN, P. G., 709
 PAULUS, A. O., (930)
 PAVGI, M. S., 119
 PEACHER, P. H., (862)
 PEEPLES, J. L., Jr., 770
 PENDERY, W., (793)
 PÉREZ, J. ENRIQUE, 569
 PETERSON, GLENN W., (618), 659
 PETERSON, ROGER S., (741), 744
 PETTIT, ROBERT E., 753
 PHELPS, WILLIAM R., 224, 736
 PIERCE, WALTER H., (238)
 PIERSON, CHARLES F., 885, 913
 PINCKARD, J. A., 172, 254
 PINE, T. S., 64, 289
 PIRONE, THOMAS P., (325)
 PITRE, H. N., 409, 570
 PITTS, RALPH, 238
 PLAUGHER, NOAH H., (22)
 POWELL, J. M., 144
 POWELSON, R. L., 806
 POWERS, H. R., Jr., 432
 PRASAD, KRISHNA, (661)
 PRASAD, N., (385), (921)
 PROVVIDENTI, R., (337)
 PURDY, LAURENCE H., 205
- RAABE, ROBERT D., 411, 519, 709
 RAHMAN, M. ABDEL, (761)
 RANGARAJAN, M., (701)
 RASKI, D. J., (27)
 RAVENSCROFT, A. V., (930)
 RAYCHAUDHURI, S. P., (88)
 REEVES, E. L., (787)
 REID, C. P. P., 345
 REYES, LUCAS, (840)
 REYNOLDS, HAROLD W., (149), 512
 RICH, A. E., (40)
 RICH, SAUL, 721
 RICO, GABRIEL, (101)
 RIFFLE, JERRY W., 126, 677
 RIGGS, R. D., 15
 RIRIE, D., (327)
 RIVERA, C. T., 453
 ROBAYO, GABRIEL, (101)
 ROBERTS, B. J., (332), 571, (631)
 ROGERS, W. E., (569)
 ROHDE, R. A., (489)
 ROHRBACH, K. G., 773
 ROMIG, R. W., (571)
 RONCADORI, R. W., (432)
 ROSENBERG, DAN Y., (768)
 ROSENKRANZ, E. E., (409)
 ROSS, ARNOLD, (736)
 ROSS, ELTON W., 527
 ROTEM, J., 635
 ROTHMAN, PAUL G., 692
 ROWELL, J. B., 800
 RUPPEL, E. G., 484
- SAINT-LAURENT, M., (356)
 SAMBORSKI, D. J., (750)
 SASSER, J. N., 664
 SCHAFFER, JOHN F., (108)
 SCHEFFER, R. P., (190)
 SCHENCK, N. C., 327, 602
 SCHIEBER, EUGENIO, 694
 SCHIFFMANN-NADEL, MINA, 251,
 867
 SCHNATHORST, W. C., 168
 SCHNEIDER, BERNARD A., 934
 SCHNEIDER, C. L., (457)
 SCHNEIDER, HENRY, (270)
 SCHOENEWEISS, DONALD F., 13, 196,
 446
 SCHREIBER, LAWRENCE R., 122,
 (573), 808
 SCHROEDER, W. T., 337
 SCHUSTER, M. L., (29)
 SCIARONI, R. H., (133)
 SECHLER, DALE T., (292), (906)
 SEELY, C. I., (852)
 SEHGAL, O. P., 862
 SHANER, G. E., (806)
 SHANKS, CARL H., Jr., 586
 SHAUGHNESSY, LYLE, (29)
 SHAW, C. G., (399)
 SHAY, J. R., (461)
 SHEPHERD, R. J., (793)
- QUIAOIT, AVELINA R., 613
 QUICK, CLARENCE R., 550

- SHER, S. A., (581)
 SILL, W. H., Jr., 11, (234)
 SIMONS, M. D., (935)
 SIMPSON, W. R., 215
 SINCLAIR, J. B., (350), (681), (686)
 SINCLAIR, W. A., 406
 SINGH, R. D., 385
 SINGH, R. S., 127, 668
 SITARAMAIAH, K., (668)
 SITTON, B. G., (54)
 SKOOG, H. A., 627
 SLEETH, BAILEY, 460
 SMERLIS, E., 356
 SMILEY, J. H., (218)
 SMITH, C. H., (650)
 SMITH, M. A., (283)
 SMITH, N. A., 190
 SMITH, RICHARD S., Jr., 181, 245,
 249, (439), 832
 SNOW, G. A., 388
 SORENSEN, E. L., (731)
 SOUSA, A. A., (424)
 SOWELL, GROVER, Jr., 661
 SPRINGER, J. K., (715)
 SPURR, HARVEY W., Jr., 424
 STALEY, JOHN M., (280)
 STAMBAUGH, WILLIAM J., (589)
 STANDIFER, LEON C., (172)
 STAPLES, ROBERT, (951)
 STEGALL, W. A., Jr., 476
 STEWART, D. M., (312), 332
 STIPES, R. JAY, 573, (808)
 STOKES, D. E., 822
 STONE, WILLIAM J. H., 414, 629
 STRETCH, A. W., (346), 715
 STUTEVILLE, D. L., 731
 SUTHERLAND, J. R., 545
 SUTTON, B. C., (208)
- TARRY, JERRY C., (181), 399
 TAYLOR, CARLTON F., (6), 701
 THAMES, WALTER H., Jr., (869)
 THOMAS, C. A., (35)
 THOMPSON, D. L., (655)
 THOMPSON, MARY RUTH, (65), (779)
 THORNBERRY, H. H., 65, (779)
 THYR, B. D., (96)
 TIWARI, V. P., (619)
- TODD, F. A., (308)
 TOLER, R. W., 316
 TOOLE, E. RICHARD, 160, 552, 945
 TORIO, JOYCE C., 346
 TOWERS, BARRY, 747
 TRAYLOR, JACK A., (278)
 TROLL, J., 489
 TROUTMAN, JOSEPH L., (139)
 TRUE, R. P., (545)
 TSAO, PAMELA W., 270
 TYSON, I. H., (62)
- VAGO, C., (764)
 VALBUENA, LUIS A., (101)
 VALCARCE, ARLAND C., (446)
 van der ZWET, T., 54, 717
 Van GUNDY, S. D., (581)
 VARNEY, E. H., (351)
 von SCHMELING, B., 645
- WAGNON, H. KEITH, 278
 WATSON, ALICE J., (348)
 WATSON, R. D., (469), (852)
 WEBB, R. E., 49
 WEBER, RAY, (224)
 WEBSTER, ORRIN J., (606), (641),
 (828)
 WEHLBURG, C., 354, 938
 WEIDENSAUL, T. CRAIG, 22
 WEINER, A., 27
 WERNHAM, C. C., 6, (363), (814)
 WESTER, R. E., (316)
 WILLIAMS, HAROLD E., (278)
 WILLIAMS, PAUL H., 192
 WILLIAMS, W. T., (800)
 WILSON, E. E., 182
 WOLFENBARGER, D. O., 908
 WORLEY, R. E., 443
 WRIGHT, W. R., 283
 WYSONG, DAVID S., 618
- YANG, SHAW-MING, 578
 YARWOOD, C. E., 639, 791
- ZAIGER, D., 892
 ZENTMYER, G. A., (892)
 ZIMMERMAN, L. H., (35)
 ZUCKERMAN, B. M., (584)

SUBJECT INDEX

- Abies* sp.: previously undescribed
 branch and stem canker with severe
 resinosis on balsam fir in Anticosti
 Island, 722
 --- bracteata: 181
Acanthorhynchus vaccinii: 539
Acer saccharum: circumstantial evi-
 dence suggests that nematodes of
 several genera may contribute to
 incidence of maple blight and die-
 back in Wisconsin, 677
Aceria tulipae: 819; 917; 951
Adenostema fasciculatum: possible res-
 ervoir of a virus which resembles
 tobacco streak, recovered from
 dodder growing on this shrub in
 southern California, 479
 Alabama: 885

- Alaska: 69; 921
- Albizia julibrissin: 945
- Alfalfa: damping-off, effects of 4 fungicides on seedlings and on controlling *Rhizoctonia solani* in sand, 435; leaf spot and damping-off distribution in Kansas, 731; *Phytophthora* root rot found in Canada for first time, potential of disease and comparison of Canadian and American isolates, 451; residue as a soil amendment for control of onion bulb rotting, 469; survey of fungi associated with wilts and root rots and an assess of damage to alfalfa in Ontario, 695
- Almond (see *Prunus*)
- Alternaria brassicicola*: 764
- dauci: 594
- ricini: 35
- tenuis: causes black core rot of mandarin oranges, 127
- Amendments: alfalfa residue, to reduce onion infection and bulb rotting, 469; soil, in India the addition of oil-cakes of margosa, castor, and peanut considerably reduced root-knot nematode populations in soil cropped to okra and tomato, 669
- Ampelopsis aconitifolia*: new host for *Xiphinema* index, 27
- Anguina* spp.: 45
- Announcements: 280; 624
- Anthrachnose: of Alpine currant, variation in *Pseudopeziza ribis* isolates from different hosts, 196
- : of cucumber, control, 340
- : of *Dioscorea alata* [winged yam], efficacy of different fungicides for control, 385
- : on Easter lily blossoms, caused by black-scale *Colletotrichum*, 751
- : new disease: of Barbados cherry in India, 518; of *Phyllanthus acidus* caused by *Colletotrichum* in India, 313
- : spotted, of hybrid tea rose petals, possibly first report of *Sphaceloma* on petals of *Rosa*, 934
- : stem, of lima bean in the South, 1965, 316
- Antibiotics: of 4 tested in combination with dimethyl sulfoxide, none gave satisfactory control of Dutch elm disease, 573; effects of several on oak wilt organism in the field, 736
- : agricultural streptomycin (100 ppm) useful spray for tomato bacterial spot disease in south Florida, 699
- : cycloheximide and phytoactin: not effective in controlling blister rust cankers in eastern white pine, 224; evaluated for control of western gall rust and aspen cankers on lodgepole and ponderosa pines, 741
- : Pentaene G8 completely inhibits seedborne fungi of chili and reduces stem rust pustules on wheat, 299
- : semicarbazone and methyl hydrazine derivatives of cycloheximide evaluated for control of comandra rust on lodgepole pine, 528
- Aphanomyces euteiches*: 91
- Aphelenchoides besseyi*: 689
- Aphis gossypii*: 305
- Apple: growth of newly transplanted trees affected in many ways by chemical soil treatments and plastic mulches used at planting time, 201; lenticel injury resulted on Golden Delicious apples stored in molded paper pulp trays and on fruit covered with filter paper both containing urea-formaldehyde wet-strength resins, but not in those containing polyamide resins, 885
- : canker: a canker disease of red delicious in Guatemala resembles blister bark disease, 694; crater canker, bark disease of Golden Delicious trees, cause unknown, extent of decline is proportional to amount of trunk injury, 3; European canker in California, importance of ascospores and macroconidia, susceptibility of leaf scars, and of rain, 182
- : nematodes, survey of orchards in Hudson Valley, 1956-1965, 783
- : powdery mildew, control by 2 new sulfites, 645
- : rots: blue-mold rot, fungicides for control, 913; collar rot of 1 to 7 year old trees propagated on East Malling II and Malling Merton 104 rootstocks, all growing in one block, 267; fruit rot caused by *Phialophora malorum*, first report in Pennsylvania, 701; perennial canker rot of, in West Virginia and Pennsylvania, 348
- : severe non-parasitic russetting and dimpling possibly result of frost injury or low temperatures during early fruit development, 151
- : stem-cavity browning associated with brown core of stored McIntosh, and possibly both disorders have common cause, 178
- : viruses: transmission studies with selective *Malus* indicators, 40; virus indexing procedures to be followed by the IR-2 Pome Fruit

- Virus Subproject, under auspices of American Phytopathological Society, 461
- Apricot (see also *Prunus*): bacterial spot, variability in leaf and fruit resistance among 17 varieties and 12 selections indicates feasibility of breeding for resistance, 112
- : limb dieback, perithecial stage of *Eutypa armeniacae* is only important source of field infection, 53
- : ring pox virus, 91 foreign varieties found to be hosts, 289
- Arceuthobium campylopodum* f. *tsugensis*: 921
- Archostaphylos manzanita*: 181
- *viscida*: 181
- Arizona: 49; 145; 414; 512; 629
- Arkansas: 862
- Armillaria mellea*: associated with dieback of *Ceanothus* spp. in Pacific Northwest, 399
- Artemisia tridentata*: 181
- Aspergillus niger*: 175; causes new sorghum stalk rot, 352
- Australia: 323; 494
- Azalea: *Rhizoctonia* leaf blight, report on disease and control, 354
- Bacteria: highly virulent *Erwinia* sp. was isolated from naturally infected lettuce and rapini plants and potato plants and tubers in Arizona, 414; method for bleaching leaves for microscope investigation of microflora on the leaf surface, 493; partial vacuum method of inoculating bean seeds with 3 common pathogens is rapid and efficient, and provides uniformly infected seed, 110; *Pseudomonas glycinea*, identification of 7 races on soybean, 557; serological relationships of races 1, 2, and 3 of *Pseudomonas solanacearum*, 275
- Bacterial blight (see Blight)
- canker: of tomato, serological techniques aid in detection of, 96
- rot (see Rot)
- spot: of apricot, variability in leaf and fruit resistance among apricots tested indicates feasibility of breeding for resistance, 112; of tomato, summary report over past 10 years on field evaluation and observations on seasonal occurrence, sources of primary inoculum, dissemination, and control in south Florida, 699
- Bamboo: rust, first report of *Uredo ignava* in the United States, found in Florida, 112
- Barley: loose smut, larger seeds produce drastically lower percentage of infected plants, 101
- : pathogenicity of *Pythium aphanidermatum*, 292
- : yellow dwarf virus, resistance in selected barley varieties, 380
- Bean: asexual sporulation of *Cephalosporium gregatum* on autoclaved Golden Bantam sweet corn medium for culturing on Pinto 111 bean, 33; partial vacuum used to inoculate seeds with pathogenic bacteria, rapid and efficient method to supply uniformly infected seed, 110; survey of lima bean diseases in the South in 1965, 316
- : angular leaf spot associated with severe defoliation of red kidney beans in Pennsylvania, 494
- : bacterial blights: in Colorado, 10% loss in 1965, 371; inheritance of reaction to halo and common blight bacteria in Nebraska #1 sel. 27 dry bean line, 29; halo blight, quick laboratory test to determine pathogenicity of *Pseudomonas phaseolicola*, 238
- : capsule mold and capsule drop of castorbean, symptoms, losses, and varietal resistance, 35
- : downy mildew, effect of low post-inoculation temperature on colonization of lima bean stems by *Phytophthora phaseoli*, 734
- : powdery mildew: control by 2 new sulfites, 645; test of fungicides on snap beans, captan + tribasic copper sulfate gave excellent control without phytotoxicity, 847
- : root rot, identity and relative importance of major root rot pathogens of snap beans in Wisconsin, 578
- : soybean cyst nematode, reproduction on 12 hosts compared, 608
- : viruses: sources of virus resistance to bean yellow mosaic, alfalfa mosaic, western ringspot, and common bean mosaic in cultivars and breeding lines, 532; yellow mosaic virus, common pea mosaic virus strongly indicated as strain of, 337
- Beet: (or tablebeet), evaluation tests in New York indicated that *Rhizoctonia* damping-off of seedlings was effectively controlled when PCNB was used in combination with other standard fungicides used to control the more important *Pythium* damping-off diseases, 614

- Berry speckle: of cranberry, common but apparently undescribed, in Wisconsin, 539
- Bioassay: possible systemic nature of six 8-quinolinol compounds tested by branch-injection into prune trees for *Cytospora fungi* control, 773
- Biological control: in India the use of soil amendments may be practical substitute for expensive chemicals for lowering root-knot nematode populations in soil, 668; study of nematode-trapping fungi and predaceous nematodes occurring in Massachusetts cranberry soils with a view toward possible biological control of plant-parasitic nematodes, 584
- Bipolaris sorokiniana-Fusarium nivale sequence in roots and crowns of winter wheat in Washington, 154
- Birdsfoot trefoil: *Leptodiscus terrestris*, first report from Missouri as root-inhabitant fungus of a legume, 753; *Thielaviopsis basicola* development enhanced by temperature stress on roots in greenhouse-grown plants, 825
- Blight: of Arizona cypress, new blight found in Georgia and South Carolina possibly caused by *Monochaetia unicornis*, 482; of guar, 9 fungi-cides tested for efficacy in control, 921; of maple, circumstantial evidence suggests that nematodes may contribute to incidence, 677
- : *Alternaria*, combined control of blight and powdery mildew of carrots, 594
- : bacterial: of beans in Colorado, 10% loss in 1965, 371; inheritance of reaction to halo and common blight bacteria in a bean cross, 29; of cotton, tests for races of *Xanthomonas malvacearum* from Australia yielded 13 infected varieties, 2 of which were resistant in the United States, 323; of radish leaf, highly pathogenic strain of *Pseudomonas maculicola* possibly resulted from repeated cropping on same ground, 192; of soybean, 16 methods tested for determining pathogenic races of pathogen, 560; of soybean, first report of identification of 7 pathogenic races, 557; *Stemphylium* and *Xanthomonas* blights, severity on tomato foliage affected by mode of irrigation in Israel, 635
- : boll, of flax, serious reduction of crop value in Texas, 840
- : brown spot needle, first report on jack pine, also found on lodgepole pine, Canada, 208
- : fire, of Stewart strain of Bartlett pear, further tests on susceptibility, 787
- : flower, of camellia, spread in Louisiana, 525
- : *Fusarium*, of turfgrasses, 942
- : gummy stem, of cantaloup, 1161 introductions screened for resistance, PI 140471 approaches immunity under field conditions, 661
- : halo: on bean, quick laboratory pathogenicity test, 238
- : leaf: of azalea, 354; northern leaf, factors affecting survival of *Helminthosporium turcicum* in corn leaf tissues, 377; *Phytophthora* leaf, of cacao seedlings, effect of adding DMSO to fungicides for control, 422
- : Rosellinia needle, on Douglas-fir seedlings, first report of *R. herpotrichioides* in California, 249
- : seedling, of cotton, use of deeply incorporated trifluralin herbicide predisposed plants to damping-off, PCNB for control, 172
- : southern: of peanut, PCNB effective control in irrigated fields, 855; of tomato, losses reduced by deep plowing, 443
- Blossom blast: of cranberry, caused by *Glomerella cingulata*, resulted in reduced fruit set, 346
- Botryosphaeria ribis*: 122
- Botryotinia ricini*: 35
- Botrytis allii*: occurrence in yellow Gilboa onions which are being grown for seed production in Israel, 21
- cinerea: testing thermal dusting with tetrachloroisophthalonitrile for control on geranium, 777
- Breadfruit: a new lethal disease in the Pacific islands can kill all branches to the main trunk in 6 weeks on a robust, full bearing tree, 892
- Bremia lactucae*: 460
- Broomrape: false, field observations on the response of Maryland tobacco to, first identified in southern Maryland in 1965, 627
- Brown core and stem cavity browning: association on stored McIntosh apples, possibly both disorders have a common cause, 178
- Brown patch disease: of St. Augustine grass, screening for resistance, 622
- Brown rot: of oranges and grapefruit: in 3-year Israel experiments the rot-causing fungus was best con-

- trolled when hot water was applied in groves 3 or 4 days after inoculation of the fruit, 867
- Buckthorn: crown rust, 1000 isolates tested for pathogenicity on key differential oat varieties, 1961-1965, 935
- Cacao: gall-inducing capacity of *Fusarium rigidiuscula* inactivated by interference from another *Fusarium* clone, 219; the roll of *Fusarium roseum* in inducing greenpoint cushion gall in the Caribbean and Latin America, 229; *Phytophthora* leaf blight, effect of adding DMSO to fungicides for control on seedlings, 422
- California: 53; 168; 181; 182; 211; 245; 249; 283; 318; 327; 439; 479; 581; 768; 832
- Calonectria rigidiuscula*: 219
- Camellia: flower blight, spread in Louisiana, 525
- Canada: 62; 144; 208; 451; 695; 722; 750
- Canker: a canker disease of red delicious apple in Guatemala, 694; previously undescribed branch and stem canker with severe resinosis on balsam fir in Anticosti Island, 722; sporophores of *Polyporus tomentosus* var. *circinatus* associated with basal cankers caused by fusiform rust fungus, 527
- : aspen, antibiotic tests for control on lodgepole and ponderosa pines, 741
- : *Corynebacterium michiganense*, diagnosis of the disease in tomato, 96
- : *Cytospora*, on honey locust, first report on thornless species, 13
- : European, development in California apple district, 182
- : pitch, fruiting of *Fusarium lateritium pini* on naturally infected pine, 476
- Canna: mosaic disease caused by cucumber mosaic virus, 1st report of natural infection of canna by CMV in the United States, 899
- Capsule mold and capsule drop: of castorbean, symptoms, losses, and varietal resistance, 35
- Carrot: 681; combined control of *Alternaria* blight and powdery mildew, 594
- Cassia: bioassay plant in method for detecting flexuous-rod virus in *Cattleya* leaf sap, 779
- Castilleja miniata: uredial and telial stages of white spored *Peridermium stalactiforme* found on, Alberta, Canada, 144
- Castor bean: influence of cropping to castor beans on populations of root-knot nematodes in soil, 132
- Cattleya*: infectious blossom necrosis, method for detecting flexuous-rod virus in leaf sap, 779
- Ceanothus* spp.: association of *Armillaria mellea* with severe dieback in Pacific Northwest, freeze of November 1955 may be predisposing factor, 399
- Celery: ozone damage in Connecticut, 721
- Cenangium singulare*: 741
- Cephalosporium* spp.: 722; associated with a decline of oak in Texas, 75
- gregatum: 598; asexual sporulation on autoclaved Golden Bantam sweet corn medium for culturing on Pinto 111 bean, 33
- maydis: 401
- stripe: of wheat, first published report of *Cephalosporium gramineum* in Michigan, 190
- Ceratocystis fagacearum*: 45; 736; 862
- fimbriata: 741
- ulmi: 473; 573; 618; 808
- Cercis canadensis*: 945
- Cercospora aleuritidis*: imperfect stage of angular leaf spot of tung, 54
- beticola: 457
- gigantea: 357
- kikuchii: 910; cause of purple stain of soybean seedcoats, 464
- Cercospora herpotrichoides*: 852
- Chaetosiphon fragaefolii*: 586
- Chemicals: chemo- and thermotherapy for control of leaf curl virus of Malvaceous plants, 88; common names for pesticides, 725; effects of several on oak wilt organism in the field, 736; 50 compounds tested did not inactivate crinkle strawberry virus when sprayed on foliage of infected plants, 843; sensitivity tests with fungicides for control of cotton seedling disease pathogens, 301; soil treatments and plastic mulches affect development of new apple trees in many ways when used at planting time, 201; two new sulfite acaricides effective in controlling apple and bean powdery mildews, 645
- : creosote, urea, and liquid borate equally effective in preventing stump surface infection by *Fomes annosus*

- after 10 months, 22
- : 2,3-dihydro-5-carboxanilido-6-methyl-1,4-oxathiin-4,4-dioxide, effective seed treatment for systemic control of stripe rust of wheat seedlings, 806
- : dimethyl sulfoxide, effect when added to fungicides for control of *Phytophthora palmivora* on cacao seedlings, 422
- : fluorescent Phosphor 2282 used to measure fungicide deposition and retention on watermelon foliage, 327
- : Freers' Dutch Elm Arrester (X-D. E. A.) tested for Dutch elm control, inconclusive results, 573
- : gibberellic acid sprays on sour cherry yellows virus infected trees increased fruit spur development at certain growth stages and environmental conditions, 240
- : growth regulators, effect of host growth inhibition by Cycocel on leaf rust infection types, 108
- : methionine and Dexon tested for control of pea root rot, divergence between results from field and greenhouse tests, 91
- : PCNB counteracts damping-off of cotton seedlings, which is a side effect of deeply incorporated trifluralin herbicide, 172
- : SMDC, DBCP, and DD used to control soil nematodes before isolation of soil fungi in a study of root rot of red pine seedling nursery, 545
- : tin compounds: fungicidal effect against 2 leaf diseases of carrot which so far could only be controlled by compounds of different chemical composition specific for each disease, 594
- : triphenyl tin combinations effective in control of tar spot of citrus, 357
- : urea-formaldehyde wet-strength resins caused lenticel injury to Delicious apples in storage in Washington, while polyamide resins did not, 885
- Cherry (see also *Prunus*): Barbados- (*Malpighia glabra*): new spot anthracnose in India, possibly caused by *Colletotrichum* sp., 518
- : viruses: bud transmission of dwarfing virus, 367; severe fruit necrosis caused by strains of green ring mottle virus, 59; Shiro-fugen as indicator for *Prunus* ringspot virus complex, 278; sour cherry yellows virus, gibberellic acid sprays increased fruit spur development of infected trees at certain growth stages and environmental conditions, 240
- Chile: 280
- Chili (see Pepper)
- Chokecherry: western X-disease virus widely distributed in North and South Dakota and Nebraska, threatens this shrub commonly used in windbreaks and as important component of wild-life plantings, 659
- Chromatography: paper, of leaf tissue, direct method, 390
- Chrysanthemum: wilt of, susceptibility of several cultivars to the Yellow Delaware Fusarium and effect of N nutrition on susceptibility, 882
- Cicca acida (see *Phyllanthus acidus*)
- Cissus rhombifolia: immune to *Xiphinema* index, 27
- Citrus spp.: 681; storage life of oranges was extended by gamma radiation of fruit inoculated after picking with *Penicillium digitatum* and *P. italicum*, Israel, 874
- : dieback and gumming of 'Robinson' tangerine twigs caused by *Diplodia natalensis* in Florida, 429
- : nematodes: citrus-root nematode found in many sweet orange and sour lemon groves in Lebanon, 960; *Tylenchulus semipenetrans* (citrus nematode) found in Trinidad for first time, 478
- : *Phytophthora citrophthora* incubation period studied in inoculated trees under natural conditions, 251
- : postharvest decay: Miller's technique for impregnating wrappers with antifungal compounds, 782
- : rots: black core rot of mandarin oranges caused by *Alternaria tenuis*, 127; brown rot, hot water treatment for control on oranges and grapefruits is most effective when applied within 3 or 4 days after inoculation, Israel, 867; brown rot syndrome in Western Australian citrus orchards, 3
- Phytophthora* spp. associated with, 494; stem-end rot fungi in Florida lemons for fresh fruit market, 860
- : tar spot, chemical control, 357
- : viruses: Arizona's Cooperative Citrus Registration-Certification Program, organization, historical development, and progress, purpose = establishment of virus-free budwood, 145; psorosis virus, transmitted through seeds of at least one citrus hybrid ('Carrizo' citrange) at levels of 15 to 31%, 81

- Cladosporium sp.: 35
 --- fulvum: 686
 Clathrus ruber: 840
 Clover: subterranean, occurrence of
 wart disease in California believed
 to be first report on this host in the
 United States, 797
 Club root: in combination with sugarbeet
 and cabbage root nematodes on
 Brussels sprouts, control by combi-
 nation soil treatments, 133
 Colletotrichum spp.: 313; 518
 --- capsici: 299
 --- --- f. cyamopsicola: 921
 --- gloeosporioides: 385
 --- graminicolum: of sorghum, spread
 in dead stalk tissue cells, and inhi-
 bition by living stalk tissue cells, 287
 --- lagenarium: 340
 --- lilii: 751
 --- orbiculare: sexual expression in, 871
 --- truncatum: 316
 Colonization: of birdsfoot trefoil roots,
 by several fungal species, first re-
 port in Missouri for Leptodiscus
 terrestris, 753; intercellular, by 3
 Malayan Fusarium isolates in Vigna
 and Hibiscus, 116
 Colorado: 309; 371; 565
 Common names: for pesticides, 725
 Compatibility: 48 pesticide-nutrient
 mixtures compared in a 3-year field
 test on watermelon in Florida, 602
 Conifers: distribution of Fomes annosus
 in natural forests of California, 832;
 16 species inoculated with 7 isolates
 of Phytophthora cinnamomi from
 various hosts and locations provides
 evidence for existence of biotypes
 of the fungus, 647
 Connecticut: 721
 Coriander: new host for Erwinia aroideae
 (soft rot), 701
 Corn: diseases in Kentucky in 1965, 218;
 sterilized Golden Bantam sweet,
 medium for asexual sporulation of
 Cephalosporium gregatum for cul-
 turing on Pinto 111 bean, 33
 ---: ergot conidia collected from sorghum
 will infect corn, 828
 ---: northern leaf blight, factors affecting
 survival of Helminthosporium turci-
 cum in leaf tissues, 377
 ---: rots: inoculation and rating proce-
 dures for resistance of corn to stalk
 rot under conditions in the South, 655;
 reaction of corn inbred lines, vari-
 eties, and hybrids to 4 fungi causing
 stalk rots in Egypt, 401
 ---: rust, morphological evaluation of 2
 pathogenic clonal lines of Puccinia
 sorghii and their inbred recombi-
 nants on a homozygous susceptible
 corn, 119
 ---: smuts: increase in susceptibility
 of inbred W22 by adding potassium to
 soil associated with decrease in
 reducing sugar and increase in nitro-
 gen in the pith tissue, 12; head smut
 in Idaho, tests indicate fungus is not
 seedborne, 215
 ---: viruses: distribution and identifica-
 tion of mechanically transmissible
 viruses in Louisiana, 325; evidence
 that corn stunt virus may not occur
 in California, 318; occurrence of
 mechanically transmissible virus of
 corn in Mississippi, 409; sugarcane
 mosaic virus, reaction of 180 sweet
 and dent varieties to Johnson grass
 strain of, 793
 ---: ---: maize dwarf mosaic: history
 and distribution in Tennessee, 427;
 host range, properties, and partial
 purification of a Missouri isolate,
 862; found in Iowa for first time, 88;
 presence established in several
 counties in Pennsylvania in 1965, 6;
 a heavy-duty mobile air blast inocu-
 lator has been designed, built, tested,
 and proved to be rapid and efficient
 for both field and greenhouse use,
 363
 ---: ---: wheat streak mosaic: host
 range and spread of virus and vector
 in South Dakota, 1962-66, 819; tests
 indicate that Ohio 3A virus culture is
 strain of wheat streak mosaic virus,
 occurrence in Ohio suggests that
 wheat and corn could be year-round
 hosts for virus and wheat curl mites
 throughout corn belt and eastern
 Great Plains, 953
 Corrections: 280; 536; 624
 Corticium fuciforme: 525
 Corynebacterium flaccumfaciens: 110
 --- michiganense: 96
 Cotton: reduction in yield caused by dis-
 eases in 1965, 350
 ---: angular leaf spot, eradicated from
 California through sanitation, 168
 ---: bacterial blight, tests for races of
 Xanthomonas malvacearum from
 Australia yielded 13 infected vari-
 eties, 2 of which were resistant in
 the United States, 323
 ---: boll rots, method for estimating
 losses, 254
 ---: nematodes: plant-parasitic nema-
 todes, seriousness and scope of
 infestation in Lower Rio Grande
 Valley of Texas, 149; root-knot

- nematode, testing experimental nematocides on 'Deltapine Smoothleaf' under Arizona conditions, 512
- : *Rhizoctonia solani* isolated from seed that were acid delinted and surface sterilized with alcoholic mercuric-chloride, 770
- : seedling damping-off, a side effect of deeply incorporated trifluralin herbicide counteracted by PCNB, 172
- : seedling disease complex variability in New Mexico, in relation to chemical control, 301
- : *Verticillium* wilt, comparison of virulence of isolates of *Verticillium albo-atrum* from several cotton-growing regions in the United States, Mexico and Peru, 930
- Cranberry: application of pesticides with "ground effect" machine, 715; study of nematophagous fungi and predaceous nematodes associated with cranberry soils in Massachusetts, 584
- : berry speckle, in Wisconsin, evaluation of fungicides and spray schedules for control, 539
- : *Glomerella cingulata* infection associated with increased blossom blast and decreased fruit set; fruit rot symptom expression also dependent upon high or low N, P, K, or S levels, 346
- : plant-parasitic nematodes, in Wisconsin, 957
- : ringspot disease, virus-like ring-spot symptoms on fruits and leaves of Searles and Howes varieties in Wisconsin, 543
- Crataegus oxyacantha*: *Gymnosporangium confusum* on, in California, 768
- Crater canker: of Golden Delicious apple, bark disease of unknown cause, extent of tree decline is proportional to amount of trunk injury, 3
- Cronartium comandrae*: 528
- *comptoniae*: 419; 744
- *fusiforme*: 388; 527; development of pycnia and aecia in albino form on inoculated pines, 842
- *ribicola*: 554
- Crucifers: new hosts for *Meloidogyne hapla*, 523
- : bacterial leaf blight: of radish, highly pathogenic strain of *Pseudomonas maculicola* possibly resulted from repeated cropping on same ground, 192
- : cabbage leaf spot, transmission of by slugs in France, 764
- : nematodes: sugarbeet and cabbage root nematodes occurring with club root organism on Brussels sprouts, marked yield increase with control, 133; sugarbeet nematodes associated with reduced yields and quality in cauliflower, cabbage, broccoli, Swiss chard and mustard, 611
- Cucumber (see Cucurbits)
- Cucurbits: decay of melons from California and Mexico on the New York market, caused by *Geotrichum candidum*, 222; viruses of, on the west coast of Mexico, 947
- : cantaloup: gummy stem blight, 1161 introductions screened for resistance, PI 140471 approaches immunity under field conditions, 661
- : cucumber: 681; anthracnose, downy mildew, and soil rot, control, 340
- : muskmelon necrotic fleck virus, new strain of tobacco mosaic virus isolated from muskmelon in Arizona, 49
- : squash: 681
- : watermelon: 3-year compatibility field tests among fungicide-insecticide-nutrient sprays in Florida, 602; surfactant influence on deposition and retention of fungicides on foliage, 327; mosaic virus, aluminum foil tested for repellency to melon aphids, 305
- Cultures: of *Cephalosporium gregatum* on Pinto 111 bean, autoclaved Golden Bantam sweet corn medium for asexual sporulation of *C. gregatum*, 33; of 4 *Geotrichum candidum* isolates on four agars in a study of plant tissue pH and susceptibility to plant and animal isolates, 681; induced sporulation of fungi on non-sterile filter paper in a moist chamber, 789; simple and inexpensive device for uniform transfer of fungus inoculum, 345; preservation of, 32 *Helminthosporium* spp. were stored under mineral oil on agar slants, 321; survival of various smut fungi cultures for 11 years under mineral oil, 800; sexual expression in *Colletotrichum orbiculare*, 871; in vitro increase in virus infectivity, 639
- Curvularia geniculata*: first report on *Sambucus* species, found on *S. chinensis*, 488
- *verruculosa*: causes severe disease of rice in India, 314
- Cuscuta cassytoides*: 593
- Cylindrocarpon* sp.: 722

- Cypress: Arizona, new blight found in Georgia and South Carolina possibly caused by *Monochaetia unicornis*, 482
- Cytospora* spp.: 773
- *chrysosperma*: 741
- *gleditschiae*: 13
- Cytosporina* sp.: perithecial stage of *Eutypa armeniacae* is only important source of apricot limb dieback field infection, not this imperfect stage, 53
- Dalbulus maidis*: 427; 570
- Damping-off: of alfalfa, caused by *Xanthomonas alfalfae*, in Kansas, 731; of alfalfa seedlings, effects of 4 fungicides on plants and on control of disease in sand, 435; detection of *Pythium* in soil, 791; of table beet, controlled by combination fungicides for *Rhizoctonia* and *Pythium* organisms, 614; of *Tephrosia vogelii*, control studies with fungicides in Puerto Rico, 484; of tomato, 124
- Decay: of western melons on the New York market, caused by *Geotrichum candidum*, 222
- Decline: of mature apple trees associated with plant-parasitic nematodes, effect of DBCP and cynem, 877; of loblolly pine, associated with *Phytophthora cinnamomi* and *Pythium* spp., 596; of oak in Texas, associated with *Cephalosporium*, 75
- Dieback: apricot limb, perithecial stage of *Eutypa armeniacae* is only important source of field infection, 53; of *Ceanothus* spp., association of *Armillaria mellea* with, 399; and gumming of 'Robinson' tangerine twigs caused by *Diplodia natalensis* in Florida, 429; of maple, circumstantial evidence suggests that nematodes may contribute to incidence, 677; and mortality, of red and jack pines caused by *Scleroderma lagerbergii* in Upper Michigan, 402
- Dieffenbachia* spp.: brown leaf spot, severe disease in Florida, proved to be caused by *Leptosphaeria* sp., 687
- Dimpling: of McIntosh and Cortland apples, possibly physiological disease, 151
- Dioscorea alata*: anthracnose of, efficacy of different fungicides for control, 385
- Diplodia natalensis*: 429; 860
- *zeae*: 655
- Diseases: check list of plant diseases previously unreported in Hawaii, 411; in Kentucky in 1965, 218; of lima beans in the South in 1965, 316; reduction in yield of cotton caused by, 350
- Distribution: amendments to records of *Cronartium comptoniae*, 744
- Dodder: parasitic on 2 principal and 3 secondary hardwood hosts each in a different family, Tanzania, 593; virus resembling tobacco streak isolated from dodder growing on a desert shrub in southern California, shrub may be reservoir for virus, 479
- Dothidella ulei*: five clones of *Hevea* sp. designated as differentials for identifying races of, 187
- Dothistroma pini*: on *Pinus radiata*, first report of this fungus in Chile, 280
- Downy mildew: of cucumber, control, 340; of lettuce, immune varieties Valverde, Calamar and Valmaine seriously affected by new race of *Bremia lactucae* in south Texas, 460; of lima bean, effect of low postinoculation temperature on colonization of stems by *Phytophthora phaseoli*, 734; of peas in western Europe, fungicidal experiments indicate that 1 spraying at the onset of flowering can reduce number of diseased pods by about half, 257; of sorghum, new sources of resistance, 641
- Drought: artificially created on loblolly pine stumps increased penetration of stumps by *Fomes annosus* 3 months after inoculation, 747
- Dutch elm disease: control studies, method for inoculation, 808; effects of SMDC on elm roots, 473; further tests with various antibiotics and fungicides gave inconclusive results; however, captan + DMSO was the best of those tested, 573; known distribution in Nebraska, 1965, 618
- Dwarf mistletoe: found on Sitka spruce in Alaska, new host record, 921
- Egypt: 401
- Elm: Dutch: effects of SMDC on elm roots, concentrations, time, and control tested, 473; further tests with various antibiotics and fungicides gave inconclusive results; however, captan + DMSO showed promise, 573; known distribution in Nebraska, 1965, 618; method for inoculation of trees with spore suspension of *Ceratocystis ulmi* for

- control studies, 808
- : phloem necrosis virus (*Morsus ulmi*) of American elm in Mississippi Delta, 751
- Enzymes: pectic, production of by Malayan isolates of *Fusarium*, 116
- Epidemiology: of bacterial spot of tomato, a study in south Florida of seasonal occurrence, sources of primary inoculum, dissemination and control, 699
- Eradication: of *Xanthomonas malvacearum* from California through sanitation, 168
- Ergot: of sorghum, host range and epidemiology in Nigeria, 828
- Erwinia sp.: highly virulent isolate from Arizona vegetables, 414
- amylovora: 787
- aroideae: 701
- carotovora: 938
- Erysiphe polygoni: 645; 847
- umbelliferarum: 594
- European canker: development in a California apple district, importance of ascospores and macroconidia, susceptibility of leaf scars, and of rain, 182
- Eutypa armeniacae: perfect stage recognized as cause of apricot limb die-back in California, not imperfect stage (*Cytosporina*), 53
- Fertilizer: application of potassium to soil greatly increased smut in corn inbred W22 by decreasing reducing sugar and increasing nitrogen in the pith tissue, 12
- Fir (see *Abies* and *Pseudotsuga*)
- Flax: rust, 16 races in varying frequencies in Canadian prairies and in North Dakota in 1963 and 1964, 62; rust, pasmo, and boll blight diseases and weathering of seed and fiber production accounted for 75 to 80% reduction in potential value of crop in Texas during 1965-66 growing season, 840
- Florida: 112; 124; 292; 602; 687; 689; 707; 822; 860
- Fomes annosus: 446; 589; 747; development of a hazard rating for, in pine stands, 510; distribution in natural forests of California, 832; found attacking and killing five previously unreported hosts, California, 181; root disease survey to determine extent of damage in natural forest stand shows that fungus is greater menace in California than past reports indicate, 439; three stump treatment chemicals effectively prevented infection by, 22
- Fraxinus americana*: graft transmission of a ringspot-like virus from, 905
- Frost: injury to wheat, 928
- Fungi: associated with wilts and root rots of alfalfa in eastern Ontario, and an assess of damage, 695; evaluation of soil microflora and isolation by apple technique in relation to the root rot disease of red pine, 545; 11 forest fungi tested for effects of cycloheximide in culture, 497; induced sporulation on filter paper, 789; inoculation technique using nylon mesh leaves fungus free of adhering growth media, can be adapted to many fungus-host combinations, 122; isolated from hypocotyls and cotyledons of fungicide-treated peanut seed grown in infested soil in Georgia, 162; Nematophagous, studied in cranberry soils in Massachusetts as possible biological control agents of phytoparasitic nematodes, 584; new records of powdery mildew fungi, 709; pepper seed microflora inhibition by antibiotic, Pentaene G8, 299; a simple and inexpensive device developed for "one-step" aseptic uniform transfer of fungus inoculum, 345; study of the important root-inhabiting fungi associated with birds-foot trefoil in Missouri, 753; stink-horn, possibly *Clathrus ruber*, unusual occurrence on golf turf, 840
- Fungicide: effect of treatment of peanut seed on soil-borne fungi at the hypocotyl and cotyledonary surfaces, 162; evaluation of fungicides and spray schedules for control of berry speckle disease of cranberry, 539; four new formulations tested for control of pecan scab for 3 seasons in Oklahoma, 599; 3-year field test of compatibility with insecticide-nutrient sprays for disease and insect control on watermelon, 602; Miller's technique for impregnating wrappers with antifungal compounds, 782; nine tested for efficacy in controlling blight of guar, 921; several evaluated for control of angular leaf spot of tung, 717
- : carbamate and copper fungicides tested for efficacy in controlling anthracnose of *Dioscorea alata* in India, 385
- : new carbamoyloxime insecticide (UC 21149, Temik) shows systemic fungicidal activity in laboratory

- studies, 424
- : cycloheximide, effects on 11 selected forest fungi in culture, 497
- : DCNA, for control of *Rhizopus* rot on peaches, 565
- : maneb, one spraying with maneb or a coordination product of zinc ion and maneb at onset of flowering can reduce number of downy mildew infected pea pods by about 50% in western Europe, 257
- : 1,4-oxathiin, 2 derivatives show chemotherapeutic activity against smut and rust diseases of grasses, 624
- : pentachloronitrobenzene: effective in controlling *Sclerotium rolfsii* when southern blight is principal disease problem in irrigated peanuts, 855; PCNB added to currently recommended table beet seed ball treatment fungicides results in the control of *Rhizoctonia solani* as well as of 2 *Pythium* species which are primarily responsible for damping-off of seedlings in New York, 614
- : Pentaene G8, see antibiotics, 299
- : several tested together: effect of various combinations of 8 on stand establishment, maintenance, and rotenoid concentration of *Tephrosia vogelii* in Puerto Rico, 484; captan + dimethyl sulfoxide was only material that showed promise of several antifungal antibiotics and fungicides tested for Dutch elm disease control, 573; captan + tribasic copper sulfate (one of several tested) gave excellent control of powdery mildew on snap beans without phytotoxicity, 847; neutral copper, agricultural streptomycin and maneb effective against bacterial spot disease of tomato in south Florida, 699; cycloheximide and phytoactin, evaluation of basal stem application of, for control of white pine blister rust on sugar pine, 554; Daconil, Dithane M-22, Dithane M-22 special, Dithane M-45, Manzate, and Manzate-D compared for control of anthracnose and downy mildew of cucumbers, 340; effect of adding DMSO to 3 copper fungicides and to malachite green for control of *Phytophthora palmivora* leaf blight on cacao seedlings, 422; nickel + dithiocarbamates, evaluation for control of wheat stem rust, 373; bioassay with six 8-quinolinol compounds for control of *Cytospora* fungi in prune trees indicates some promise of effective systemic action, 773; PCNB, Thimer, and thiram tested for control of snow mold on four varieties of bluegrass in Alaska, 69; combination with surfactants, retention of on watermelon foliage measured by a fluorescent chemical, 327; effects of thiram, captan, pentachloronitrobenzene and chloranil on alfalfa seedlings and on controlling *Rhizoctonia solani* in sand, 435
- : SMDC, effects on elm roots, concentration-time-control tests on *Ceratocystis ulmi*, 473
- : sodium-o-phenylphenate, 2-aminobutane, and thiabendazole tested for control of blue-mold rot of apples, 913
- : tetrachloroisophthalonitrile, thermal dusting for control of *Botrytis cinerea* on geranium in the greenhouse, 777
- Fusarium* spp.: 122; 695; colonization by 3 Malayan isolates in *Vigna* and *Hibiscus*, 116; toxin production by non wilt-inducing isolates, 261
- *decemcellulare*: 229
- *graminearum*: 401
- *lateritium pini*: fruiting on naturally infected pine pitch cankers in North Carolina, 476
- *moniliforme*: 401; 655
- *nivale*: 154
- *oxysporum*: 469; 882
- --- *f. apii*: 391
- --- *f. lycopersici*: 707; serological differentiation of race 1 from race 2, 444
- *rigidiuscula*: 219
- *roseum*: 26; 229; 942
- *solani*: 160; pathogenic to bean and pea, 578
- wilt (see Wilt, *Fusarium*)
- Fusicladium effusum*: 599
- Ganoderma lucidum*: 243
- Georgia: 162; 306; 393; 482; 839
- Geotrichum candidum*: 222; 3 plant and 1 human isolate of: study to test pathogenicity of and relationship between tissue pH and susceptibility of 15 puncture-wound inoculated plant tissues, 681
- Geranium: bacterial stem rot and leaf spot, satisfactory technique for checking presence of bacteria in commercial plants, 515; *Botrytis cinerea*, testing thermal dusting with tetrachloroisophthalonitrile for control, 777
- Gibbera compacta*: 539

- Gloeosporium perennans*: 348
- Glomerella cingulata*: 346
- Gloxinia*: *Thielaviopsis basicola* rot of rhizomes from Europe, 709
- Gramineae: pathogenicity of 91 morphologically similar isolates of *Helminthosporium* to 30 gramineous species, 382
- Grape: hot-water treatments effective in eliminating *Pratylenchus vulnus* from grapevine roots, however soil treatments should still be used, 858
- Grasses: citrus nematode found on an indigenous Florida grass, *Andropogon rhizomatus*, 822; *Hypsoperine graminis* in Kansas, 396; root damage resulting from HCN produced by *Marasmius oreades*, 264; 2 new systemic fungicides (derivatives of 1,4-oxathiin) show chemotherapeutic activity against rusts and smuts, 624; 44 tested for susceptibility to aster yellows virus, host range now includes 8 grasses (including oats), 17
- : Bermuda grass, new host for red thread, found in Mississippi, 525
- : bluegrass, snow mold of, three chemicals tested for control on four varieties of bluegrass in Alaska, 69
- : gamagrass: new host of *Dalbulus maidis*, 3rd plant species reported to be accepted by leafhopper for development from egg to adult, 570; new perennial host of maize dwarf mosaic virus, Illinois, 65
- : guinea, host range and epidemiology of the sorghum ergot organism in Nigeria, 828
- : Johnson grass: maize dwarf mosaic virus, positive transmission from one source, first report in Kansas in 1965, 11; 2 distinct strains of maize dwarf mosaic virus in northeastern U.S. identified by host reaction and serological relationships, 814
- : Kentucky bluegrass: evidence for existence of physiologic races of stripe smut, 430; severity of melting-out reduced by use of resistant varieties, avoiding excessive nitrogen fertilization, and raising cutting height of turf, 703
- : St. Augustine: screening for tolerance to *Rhizoctonia solani*, 622; seedlings mechanically inoculated with maize dwarf mosaic virus resulted in 80% infection, potential threat to South, 441
- : turfgrass: effects of nematocides on growth, 489; *Fusarium* blight of turfgrasses, 942; pseudo-root-knot nematode, found in Texas for first time, 645; unusual occurrence of stinkhorn fungus on golf turf, has unsightly appearance, viscous nature, offensive odor, and distorts putting surface, 840
- : turf and forage, studies of stripe smut prevalence, pathogenicity and control in New Jersey, 294
- : western wheatgrass, first record of seed-gall nematode on, found in Montana, 45
- : zoysia: first report of rust on *Zoysia* spp. in Alabama, 885; occurrence of rust in Georgia, 839; *Puccinia zoysiae* found on *Zoysia* spp. in Mississippi, 770
- Greenpoint cushion gall: of cacao, the role of *Fusarium roseum* in causing the disease, 229; of cacao, gall-inducing capacity of *Fusarium rigidiuscula* inactivated by interference from another *Fusarium* clone, 219
- Guar: blight of, 9 fungicides tested for efficacy in control, 921; tobacco ring spot virus, new occurrences in Texas and Oklahoma, 7
- Guatemala: 694
- Guignardia vaccinii*: 539
- Gymnosporangium confusum*: in California, first report in western hemisphere, 768
- Hardwoods: dodder parasitic on seedlings in Tanzania forest nursery, 593
- Hawaii: 411; 519
- Hazard rating: for *Fomes annosus* in Virginia, guidelines provided by a survey which are based on criteria of soil texture, depth of A horizon and height of water table to predict rate of spread in infected loblolly pine stands, 510
- Helminthosporium* spp.: longevity and pathogenicity of 32 species stored under mineral oil on agar slants, 321; pathogenicity of 91 morphologically similar isolates to 30 gramineous species, 382
- turcicum: factors affecting survival in corn leaf tissues, 377
- vagans: 703
- Hemicyclophora arenaria*: 581
- Hemp sesbania: comparative rates of reproduction of soybean cyst nematode on 12 hosts, 608
- Herbicides: diuron, fall applications to control weeds in winter wheat reduced incidence and severity of foot rot, 852

- : trifluralin predisposed cotton to Rhizoctonia seedling blight, PCNB may be practical control, 172
- Heterodera cruciferae: 133
- glycines: 15; 548; 608
- punctata: occurrence and distribution, 869
- schachtii: 133; 650
- tabacum: 506
- Hibiscus esculentus: effect of culture filtrates of Fusarium isolates on wilt production, 261; intercellular colonization and the production of pectic enzymes by Malayan isolates of Fusarium, 116
- rosa-sinensis: leaf curl virus, chemo- and thermotherapy for control, 88
- Honeylocust: Cytospora canker on, first report on thornless species, 13; Polyporus lucidus associated with sudden death of tree in Mississippi, 243
- Horizon, A: depth of, as it relates to loblolly pine infection by Fomes annosus, 510
- Hosts: 10 weed species collected in Mississippi suspected to be natural summer hosts for barley yellow dwarf virus, 692
- Hot water treatments: (see temperature)
- Hydrogen cyanide: production by Marasmius oreades damages cereal and grass roots, 264
- Hymenoclea salsola: serves as native host to sheath nematode which is a native to California soils, 581
- Hypoxylon pruinaum: 741
- Hypsoperine graminis: 645; in Kansas, 396
- Idaho: 215
- Ilex spp.: 4 cultivars tested for pathogenicity of Meloidogyne spp., 664
- Illinois: 65; 446
- India: 313; 314; 612; 668; 701
- Insecticides: combined with fungicides and nutrients in 3-year compatibility tests for spraying watermelon foliage in Florida, 602
- : 2-methyl-2-(methylthio)propionaldehyde O-(methylcarbamoyl)oxime, a new insecticide-acaricide, shows nematocidal and systemic fungicidal properties in the laboratory, 424
- International facility: to prevent importation of foreign plant pathogens, 803
- Interregional Research Project-2 (IR-2): Deciduous Tree Fruit Virus Project, originally designed to index viruses of stone fruits, has been extended to include indexing for latent viruses of pome fruits, particularly apples, 461
- Iowa: 88
- Irrigation: relationship between mode of irrigation and severity of tomato foliage diseases in Israel, 635
- Isariopsis griseola: 494
- Isolation: quarantine laboratory, preliminary outline of requirements needed by such a research facility (e. g. security to prevent escape) for the study of plant pathogens, viruses in particular, that have so far been excluded from entry into various countries, 803
- Israel: 21; 711; 867; 874
- Juniperus sabina: Gymnosporangium confusum on, in California, 768
- Kansas: 11; 234; 396; 731
- Kentucky: 218
- Leaf: bleaching of, for microscope investigation of microflora on leaf surface, 493; disposable leaf presses for obtaining plant sap for serodiagnosis, 822
- Leaf press: disposable, to obtain sap from plants for serodiagnosis, 822
- Leaf scald: of sugarcane, found in Puerto Rico, first report of occurrence in one of the larger islands of the Antilles, 569
- Leaf spot: angular: associated with severe defoliation of red kidney beans in Pennsylvania, 494; of cotton, Xanthomonas malvacearum eradicated from California through sanitation practices, 168; of strawberry, found in Wisconsin for first time, 167; of tung, effect on yield and oil content of fruit from trees sprayed with Bordeaux mixture or petroleum oil for 5 years, 54; of tung, evaluation of various fungicides for control, 717
- : bacterial: of alfalfa and new hosts, distribution in Kansas, 731; of geranium, inoculation technique, 515
- : brown, severe on several Dieffenbachia spp., proved to be caused by Leptosphaeria sp., 687
- : of cabbage, transmitted by slugs, 764
- : Cercospora, of peanut, technique for forecasting, 810
- : of strawberry, relative resistance of 63 varieties and selections studied at 2 locations for 2 years, 105

- Lebanon: 960
 Lecanosticta acicola: 208
 Lenticel injury: of apples in storage caused by urea-formaldehyde resins, extent of injury also dependent upon temperature in Washington, 885
 Leptodiscus terrestris: colonization of birdsfoot trefoil roots in Missouri, first report of, 753
 Leptosphaeria sp.: proved to cause brown leaf spot of Dieffenbachia, 687
 Lespedeza: comparative rates of reproduction of soybean cyst nematode on 12 hosts, 608
 Lettuce: bacterial soft rot of Iceberg (Great Lakes) in Florida Everglades, 938
 ---: calico, caused by alfalfa mosaic virus not by tobacco ringspot virus, 629
 ---: downy mildew, immune varieties Valverde, Calamar and Valmaine seriously affected by new physiologic race of Bremia lactucae in south Texas, 460
 ---: highly virulent Erwinia sp. isolated from naturally infected plants in Arizona, 414
 Libocedrus decurrens: Fomes annosus found killing trees within native range in California, 181
 Lily: Easter: anthracnose on blossoms caused by black-scale Colletotrichum, 751; root-lesion nematodes, control with phosphate pesticides, 923
 ---: Lilium regale, bulbs from the Netherlands infected with Aphelenchoides fragariae were intercepted by plant quarantine inspectors, 798
 Longidorus vineacola: 711
 Loose smut: in Funza barley, influence of seed size on incidence of disease, 101
 Lophophacidium hyperboreum: pathogenicity of, 356
 Losses: tomato bacterial spot, evaluation of causes of this important disease and effective methods to combat its development and dissemination in south Florida, 699
 Louisiana: 325; 525; 596; 686
 Macrophomina phaseoli: 352; found on 3 new hosts in Pakistan, 171; responsible for most losses in forest nurseries in California, control by soil fumigation, 245
 Macrosteles fascifrons: 17
 Malvaviscus arboreus: leaf curl virus, chemo- and thermotherapy for control, 88
 Marasmius oreades: fairy ring fungus produces HCN which damages grass roots, 264
 Maryland: 351; 627
 Massachusetts: 151; 584
 Melampsora lini: 62; 840
 Meloidogyne spp.: 126; 149; 664; 915
 --- hapla: 132; new hosts of, 523
 --- incognita: 132
 --- --- acrita: 512
 --- javanica: 132; 668
 --- thamesii: 132
 Melting-out: of Kentucky bluegrass, severity reduced by use of resistant varieties, avoiding excessive nitrogen fertilization, and raising mowing height of turf, 703
 Methods and techniques: aluminum foil tested for repellency to melon aphids in watermelon mosaic virus control, 305; apparatus for mist culture of wheat, for study of influence of stripe rust on root development, 369; apple technique employed to isolate fungi from red pine seedling bed nursery in West Va. previously treated with several nematocides, 545; bleaching leaves for microscope investigation of microflora on the leaf surface, 493; comparison of 2 methods for inoculating Nuttall oaks with heart rot fungi, 552; deep plowing reduces losses due to southern blight on tomato, and generally improves plant vigor and fruit yields, 443; detection of Pythium in soil, 791; direct method for qualitative paper chromatography of leaf tissue, 390; disposable leaf presses for obtaining plant sap for serodiagnosis, 822; effects of surfactants on deposition and retention of fungicides on watermelon foliage measured by a fluorescent chemical, 327; estimating losses from cotton boll rots, 254; for evaluating hundreds of lines and varieties of wheat at maturity for resistance to common root rot, 26; forecasting leafspot on peanuts, 810; "ground effect" machine shows promise as a vehicle for applying pesticides on cranberry, 715; a heavy-duty, rapid, mobile and efficient pressure inoculator for mechanical transmission of maize dwarf mosaic virus was designed, built, and tested for both greenhouse and field use, 363; increment core sampling of Pinus spp. for Fomes annosus, 589; for identification of Puccinia graminis, permits sanitation, positive separa-

tion, and reduction in time needed to identify components of racial mixtures, 312; induced sporulation of fungi on filter paper, simple and inexpensive method, 789; inoculating geranium cuttings with *Xanthomonas pelargonii* to determine presence of bacteria in commercial plants, 515; inoculation of elms with spore suspension of *Ceratocystis ulmi*, 808; inoculation method is most efficient and precise of 16 tested for determining pathogenic races of soybean bacterial blight pathogen, 560; inoculation technique using nylon mesh can be adapted to many fungus-host combinations, 122; in vitro increase in virus infectivity, 639; isolating *Verticillium* spp. from field soil, 897; metal leaf presses in pliers to express plant sap for serodiagnosis, 822; Miller's technique for impregnating wrappers with antifungal compounds, 782; needle weight determination method accurately estimates intensity of *Rhabdocline* needle cast of Douglas-fir, 406; partial vacuum method of inoculating bean seeds with pathogenic bacteria is rapid and efficient, and provides uniformly infected seed, 110; quick laboratory test to determine pathogenicity of *Pseudomonas phaseolicola* on bean, 238; rapid flotation-sieving technique for extracting nematodes from soil, does not vary with time of year, 954; rapid method of assaying pathogenic potential of *Puccinia graminis tritici* populations for use in epidemic prediction and in detection of new, dangerous virulences, 673; in rating procedures for evaluating corn resistance to stalk rot caused by *Diplodia zeae* and *Fusarium moniliforme*, the toothpick and syringe methods were superior to inoculator method in the South, 655; screening St. Augustine grass for tolerance to *Rhizoctonia solani*, 622; storage of pycniospores of *Puccinia graminis secalis* in liquid nitrogen, 576; serological procedures simplify differentiation of race 1 from race 2 of *Fusarium oxysporum* f. sp. *lycopersici*, 444; simple device for uniform transfer of fungus inoculum, 345; a simple, efficient, and inexpensive method of collecting strawberry leaves for virus infection indexing, 426; simple, inexpensive, continuous sampling spore trap, 906;

simple method for detection of a flexuous-rod virus associated with infectious *Cattleya* blossom necrosis, 779; storage of urediospores of *Puccinia graminis tritici* in liquid nitrogen, 502; survival of smut cultures under mineral oil for 11 years, 800; two methods for test inoculation of ponderosa pine with western gall rust, 550

Mexico: 598; 947

Michigan: 190; 402

Millet: ergot conidia collected from guinea grass will infect millet, 828

Minnesota: 557

Mississippi: 35; 243; 388; 409; 525; 570; 692; 770

Missouri: 753; 862

Mock orange: powdery mildew found for first time in Southeast (Georgia), 306

Monochaetia unicornis: 482

Montana: 45

Morus tatarica: inoculation technique to test pathogenicity of *Fusarium* spp. on, 122

Muskmelon (see Cucurbits)

Mycosphaerella aleuritidis: 717; perfect stage of angular leaf spot of tung, 54

--- *citrullina*: 661

--- *fragariae*: 105

Myrica gale: sweet fern blister rust on, 2 specimens found in Northwest Territories, 419

Myzus persicae: 613

Nebraska: 618

Nectria galligena: 182

Needle cast: *Rhabdocline*, of Douglas-fir, needle weight determination method accurately estimates intensity of disease, 406

Nematocides: effects on turfgrass growth, 489; new carbamoyloxime insecticide (UC 21149, Temik) shows nematocidal activity in laboratory studies, 424; phosphate pesticides control root-lesion nematodes in Oregon Easter lily plantings, 923; volatile and nonvolatile tested for control of tobacco cyst nematode, 507

---: 1,3-D and DD mixture, tested over a 2-year period for control of sugar-beet nematode in field plots cropped to vegetables, 611

---: DBCP: more effective than cynem when response of mature apple trees to both treatments was tested in the Hudson Valley, 877; 1,2-dibromo-3-chloropropane, 1,3-dichloropropane, and 1,2-dibromoethane tested for effect of exposure time and dose

- on nematode eradication, under controlled conditions, 915
- : *O,O*-diethyl *O*-2-pyrazinyl phosphorothioate, used as a dip eradicated soybean cyst nematodes from roots of plants in pot tests, 548
- Nematodes: in maple blight and dieback areas in Wisconsin may contribute to incidence of these diseases, 677; rapid flotation-sieving technique for extracting nematodes from soil, does not vary with time of year, 954; relation between nematocide soil treatment for control in red pine seedling bed nursery and effect on soil fungi isolated in study on cause of root rot of conifers, 545; survey of apple orchards in Hudson Valley, 1956-65, 783
- : *Aphelenchoides fragariae*, found on 3 tree peony species from Japan and on *Lilium regale* bulbs from the Netherlands by U.S. plant quarantine inspectors, 798
- : citrus: found on new grass host and associated with additional plants in two surveys, 822; found in Trinidad for first time, 478
- : citrus-root, in many sweet orange and sour lemon groves in Lebanon, 960
- : cotton root-knot, efficacy and residual nature of experimental nematocides tested for control on Deltapine cotton in Arizona, 512
- : cyst, notes on occurrence and distribution of *Heterodera punctata*, 869
- : dagger, can cause severe decline of *Vinca minor*, 420
- : ectoparasitic, disease of onion associated with *Longidorus vineicola* serious in Israel, 711
- : foliar, cause of "leaf stripe" disease of rubber plant in south Florida, 689
- : parasitic, effects of nematocides on turfgrass growth, 489
- : plant-parasitic: of cotton, percentage of infestation in Lower Rio Grande Valley of Texas, 149; on cranberries in Wisconsin, 957; response of mature apple trees to DBCP and cynem in the Hudson Valley, 877
- : predaceous, extracted from cranberry soils, identified and frequency counts made for future application to biological control of plant-parasitic species, 584
- : pseudo-root-knot, of turf grasses, found in Texas for first time, 645
- : reniform, varietal reaction of 24 sweetpotato selections to, 500
- : root-knot: effects of exposure time and nematocide dose on eradication, under controlled conditions, 915; pathogenicity of *Meloidogyne* spp. on *Ilex* cultivars, 664; *Meloidogyne* (may be undescribed species) possibly associated with decline of ponderosa pine in New Mexico, 126; new hosts for *Meloidogyne hapla*, 523; cropping to castor beans reduced *Meloidogyne hapla* and *M. thamesii*, but reproduction by *M. incognita* and *M. javanica* was excellent, 132; (*Meloidogyne javanica*) significant reduction of eggs, larvae, and mature females in soil as result of oil-cakes of margosa, castor, and peanut in India, 668
- : root-lesion: control in Oregon Easter lily plantings with phosphate pesticides, 923; hot-water treatments effective in eliminating *Pratylenchus vulnus* from grapevine roots, however soil treatment should still be used, 858
- : seed-gall, first record on western wheatgrass, and first record on grass in Montana, 45
- : sheath, a native to California soils, and *Hymenoclea salsola* serves as a native host, 581
- : soybean cyst nematode: additional weed hosts, 15; comparative rates of reproduction on 12 host plants, 608; elimination from roots of plants by chemical dips in pot tests, 548
- : sugarbeet: associated with reduced yields and quality in vegetable crops, 2-year field experiments with 1,3-D and DD mixture for control, 611; evaluating sugarbeet varieties and breeding lines for tolerance to nematode invasion, no outstanding resistance was observed, 650; possible interaction between sugarbeet and cabbage root nematodes and club root organism on Brussels sprouts, evaluation of combination soil treatments for control, 133
- : tobacco cyst, volatile and nonvolatile nematocides tested for control, 507
- : *Xiphinema* index, 5 new hosts for, 27
- New diseases: blight of Arizona cypress found in Georgia and South Carolina possibly caused by *Monochaetia unicornis*, 482; new disorder of strawberry fruit, 327; Pingalap disease of breadfruit in the Pacific islands has destroyed an estimated 80,000 trees in 2 years, it can kill

- all branches to the main trunk in 6 weeks, 892; previously undescribed branch and stem canker with severe resinosis on balsam fir in Canada, 722; sorghum stalk rot caused by *Aspergillus niger*, 352; stunt disease of peanut, undescribed virus causes severe dwarfing and fruit suppression, found in Virginia, 139
- : in India: new spot anthracnose of *Malpighia glabra* (Barbados-cherry) possibly caused by *Colletotrichum* sp., 518; anthracnose of *Phyllanthus acidus*, 313
- : in the United States: crater canker of Golden Delicious apple trees, bark disease of unknown cause, Washington, 3; ozone injury to celery in Connecticut is possibly first report in United States, 721
- New distribution: check list of plant diseases previously unreported in Hawaii, 411
- : apple fruit rot, caused by *Phialophora malorum*, in Pennsylvania, 701
- : angular leaf spot of strawberry in Wisconsin, 167
- : camellia flower blight in southwest Louisiana, 525
- : *Dothistroma pini* in Chile, found on *Pinus radiata*, 280
- : false broomrape on tobacco in Maryland, 627
- : *Fomes annosus* found on *Libocedrus decurrens* in California, 181
- : *Gymnosporangium confusum* in western hemisphere for first time, found in California, 768
- : *Lecanosticta acicola* found in Canada, 208
- : *Macrophomina phaseoli* found on *Physalis peruviana*, *Rumex dentatus* and *Solanum nigrum* in Pakistan, 171
- : maize dwarf mosaic virus on corn in Iowa, 88
- : nematodes: citrus nematode (*Tylenchulus semipenetrans*) in Trinidad, 478; pseudo-root-knot nematode in Texas, 645; seed-gall nematode on grass in Montana, 45
- : *Phytophthora cinnamomi*, on loblolly pine in Louisiana, 596
- : --- *fragariae* race A-5 found in commercial strawberry plantings in the eastern United States, 351
- : *Polyporus lucidus* (*Ganoderma lucidum*) on honeylocust in Mississippi, 243
- : --- *tomentosus* spread in slash pine plantations in the southeastern United States, 527
- : powdery mildew: on mock orange in Georgia and the Southeast, 306; new records of powdery mildew fungi, 709
- : *Puccinia zoysiae*, in Mississippi, 770
- : *Rosellinia herpotrichioides*, in California, found on Douglas-fir seedlings, 249
- : root rot, of alfalfa caused by *Phytophthora megasperma* has been found in Canada, 451
- : rust: on zoysia grass in Alabama, 885; zoysia rust in Georgia, 839
- : sugarcane leaf scald, in Puerto Rico, first report on occurrence in one of the larger islands of the Antilles, 569
- : *Uredo ignava*, first report in the United States, found on bamboo in Florida, 112
- : wart disease, on subterranean clover in the United States, 797
- New Hosts: alfalfa mosaic virus, first known report as causal agent for a disease (calico) of head lettuce in the field, Arizona, 629
- : apricot ring pox virus, 91 apricot varieties from foreign sources found to be hosts for, 289
- : aster yellows virus, on several grass hosts, 17
- : barley yellow dwarf virus, recovered from 5 weed species collected in Mississippi, believed to be natural summer hosts, 692
- : cucumber mosaic virus, causes mosaic disease of canna, 1st report of natural infection of canna by CMV in the United States, 899
- : *Curvularia geniculata*, on *Sambucus chinensis*, 488
- : --- *verruculosa*, on rice in India, 314
- : *Cytospora* canker, on thornless species of honeylocust, 13
- : *Dalbulus maidis*, on gamagrass, this is the 3rd plant species reported to be accepted by this leafhopper for development from egg to adult, 570
- : dwarfmistletoe, on Sitka spruce, found in Alaska, 921
- : *Erwinia aroideae*, on pear fruits and coriander leaves in India, 701
- : *Fomes annosus*, on *Pinus patula*, *Abies bracteata*, *Artemisia tridentata*, *Archostaphylos viscida* and *A. manzanita*, found in California, 181
- : *Lecanosticta acicola* (conidial state of *Systremma acicola*), on *Pinus banksiana*, 208

- : *Macrophomina phaseoli*, on *Solanum nigrum*, found in Pakistan, 171
- : maize dwarf mosaic, on gamagrass, Illinois, 65
- : root-knot nematode, found on ponderosa pine in New Mexico, believed to be first report, 126; for *Meloidogyne hapla*, 523
- : seed-gall nematode (*Anguina* spp.), on western wheatgrass, 45
- : soybean cyst nematode, on seven additional weed hosts in Arkansas, 15
- : *Tylenchulus semipenetrans*, on *Andropogon rhizomatus*, 822
- : *Xanthomonas alfalfae*, readily infected several in greenhouse tests, 731
- : *Xiphinema* index, on 5 new hosts, 27
- : red thread fungus, on Bermuda grass in Mississippi in 1965-66, 525
- New Jersey: 294; 351
- New Mexico: 126; 175; 301
- New York: 783; 877
- New Zealand: 730
- Nigeria: 606; 641; 828
- Nilaparvata lugens*: 453
- Nitrogen: effect of various levels in the soil on stalk rot of corn incidence, 655
- : liquid: storage of pycniospores of *Puccinia graminis secalis* in, 576; storage of urediospores of *Puccinia graminis tritici* in, 502
- North Carolina: 136; 308
- North Dakota: 62
- Nutrients: effects of varying combinations of N, P, K, and S on fruit rot symptoms in cranberry, 346; high and low N effects on susceptibility of chrysanthemum to *Fusarium* wilt, 882; nutrient factors plus tissue pH in mesocarp and epicarp of orange may be involved in preventing infection by *Geotrichum candidum*, 681
- Oak: release of aeciospores from *Cronartium fusiforme* galls on slash and loblolly pine found to correlate well with appearance of susceptible young leaves of water oak in 2 spring seasons in southern Mississippi, 388
- : decline, of live, southern red, water, and post oaks in Texas associated with *Cephalosporium*, 75
- : *Fusarium solani*, isolated from oak cankers can infect young oak stems and is more virulent than strains of the fungus from tupelo, cottonwood, and yellow-popular, 160
- : fusiform rust, 432
- : heart rot, comparison of 2 methods for experimentally inoculating Nuttall oaks, 552
- : wilt: distribution extended in Arkansas, 862; evaluation of small oak bark beetle as a vector, 45; of northern pin, field studies of antibiotics and chemicals for control, 736
- Oats: pathogenicity of *Pythium aphanidermatum*, 292
- : crown rust: adult plant resistant hybrids of CI 3034 should be searched for stem rust resistance, 631; 3000 isolates tested for pathogenicity on key differential oat varieties, 1961-1965, 935; 18 races new to North America isolated during recent surveys in Canada, 750
- : stem rust: discovery of a new and virulent race, 6AFH, in the Uniform Oat Rust Nursery, which attacks varieties with the A, B, D, E, F, and H genes for resistance, 751; new type of resistance of adult plants may be effective against all North American races, 631; physiologic races of *Puccinia graminis* in the United States in 1964, 332
- Ohio: 951
- Oidium caricae*: 519
- Oil: longevity and pathogenicity of *Helminthosporium* spp. stored under mineral oil, 321
- : spray, 1.25% and 2.85% of a light oil emulsion delayed field spread of pepper viruses but did not increase yields, 158
- Oiliopsis taurica*: 635
- Oklahoma: 7; 599
- Okra (see *Hibiscus esculentus*)
- Onion: *Botrytis allii* attacking yellow Gilboa variety which is being grown for seed production in Israel, 21; bulb rotting by soilborne pathogens, especially *Fusarium oxysporum*, greatly reduced by alfalfa residue amendment, onion residue increased rotting, 469; disease associated with ectoparasitic nematode serious in Israel, 711
- Oregon: 59
- Ornamentals: 1965 survey in Georgia nurseries to determine prevalence of root rot organisms and which fungi were involved, 393
- Ozone: injury to celery, first report in Connecticut and possibly in United States, 721

- Pacific Islands: 892
 Pakistan: 171
 Papaya: powdery mildew, susceptibility of senescent leaves of fruiting plants, Hawaii, 519; relationship of virus incidence with distance and time with-in a field, 908
 Paratylenchus curvatus: 877
 Parthenocissus tricuspidata: new host for Xiphinema index, 27
 Pasm: of flax, serious reduction of crop value in Texas, 840
 Pathogenicity: of Lophophacidium hyperboreum, 356
 Pea: common pea mosaic virus, strongly indicated as strain of bean yellow mosaic virus, 337
 ---: downy mildew, number of diseased pods reduced about 50% with 1 fungicidal spraying at the onset of flowering in western Europe, 257
 ---: root rot, study of chemical control using methionine and Dexon, divergence between results from field and greenhouse tests, 91
 Peach (see also Prunus): 681
 ---: latent viruses, transmission in commercial seed, 740
 ---: mule's ear virus disease, 64
 ---: Rhizopus rot, control studies with preharvest sprays of 2,6-dichloro-4-nitroaniline (DCNA) in combination with adjuvants and with use of both hard and soft water under Colorado conditions, 565
 Peanut: mycoflora isolations from peanut hypocotyls and senescent cotyledons reveals a similarity in the spectra of fungi harbored by tillering plants grown in 2 Georgia soils from fungicide-treated seed, 162
 ---: Aspergillus crown rot, serious outbreak in New Mexico in 1965, 175
 ---: bacterial wilt: disease reaction and symptom expression of 17 cultivars, 520; rapid detection of, in plant by agar-diffusion method, 836
 ---: leafspot, technique for forecasting, 810
 ---: southern blight, PCNB effective control of Sclerotium rolfsii when blight is principal disease problem in irrigated fields, 855
 ---: stunt, new virus disease causes severe dwarfing and fruit suppression, mechanical, graft, dodder, and aphid, transmission, 139
 ---: virus, severe stunting and yield reduction of 2 varieties and obvious spread of unidentified disease in 5 North Carolina locations, 136
 Pear: Stewart strain of Bartlett, further tests on susceptibility to fire blight, 787
 ---: decline virus, new brown leaf-vein symptom associated with infected greenhouse-grown plants, 270
 ---: soft rot, of fruit, new host for Erwinia aroideae, India, 701
 Pecan: scab, 4 new fungicide formulations tested for control in Oklahoma, 599
 Penicillium digitatum: 874
 --- expansum: 913
 --- italicum: 874
 Peniophora gigantea: 747
 Pennsylvania: 6; 348; 494; 701
 Peony: tree, 3 species from Japan intercepted by plant quarantine, buds were infected with the nematode Aphelenchoides fragariae, 798
 Pepper: 681; seed, fungal microflora completely inhibited by new antibiotic, Pentaene G8, 299
 ---: viruses: new strain of tobacco mosaic virus naturally occurring on chilli pepper variety puri orange in India, virus has wide host range, 619; oil sprays delay field spread but do not increase yields, 158
 Peppermint: Verticillium wilt of, most effective soil fumigants of several tested, controlled for 3 consecutive crops, 97
 Peridermium stalactiforme: 144
 --- harknessii: 550; 741
 Peronospora pisi: 257
 Pesticides: application on cranberry with "ground effect" machine shows promise, 715; common names for, 725
 Phialophora malorum: 701
 Phomopsis citri: 860
 Phyllactinia guttata: 306
 Phyllanthus acidus [=Cicca acid]: new anthracnose disease of, in India, 313
 Physiological diseases: lenticel injury to apples in storage in Washington, 885
 Physoderma trifolii: 796
 Phytophthora spp.: 393; 695
 --- cactorum: 267; 283
 --- cinnamomi: 596; evidence for existence of biotypes of, based on differential pathogenic effects to the same and different broadleaf and coniferous evergreens, 647; 13 isolates from various hosts and locations differed in degree of pathogenicity to 6 Taxus cultivars, 84
 --- citricola: 494
 --- citrophthora: 251; 494; 867
 --- fragariae: race A-5, found in two

- commercial strawberry plantings for first time in eastern United States, 351
- hibernalis: 494
 - infestans: 686
 - megasperma: 451
 - --- var. sojae: 104
 - palmivora: 422
 - parasitica: 686
 - phaseoli: 734
- Picea sitchensis*: first report of dwarf-mistletoe on, 921
- Pinus* spp.: annosus root rot, significantly higher incidence of infection revealed by core sampling than by field diagnosis, 589; cycloheximide and phytoactin evaluated for control of western gall rust and aspen cankers on lodgepole and ponderosa pines, 741; distribution and severity of *Fomes annosus* root rot in Illinois pine plantings, 446; fruiting of *Fusarium lateritium pini* on naturally infected pitch cankers of a Virginia pine in North Carolina, 476; fusiform rust, time of maximum teliospore production and weather conditions necessary for maximum infection in South Carolina, 432; incidence of *Polyporus tomentosus* fruiting on infected trees in a Georgia slash pine plantation, 360; loblolly pine stands given low, intermediate, or high hazard ratings for benefit of management foresters, 510; occurrence of sweet fern blister rust in Alberta and Northwest Territories could be serious threat to hard pine nurseries and plantations, 419; red pine root rot study conducted in West Virginia nursery, 545; *Scleroderris lagerbergii* responsible for widespread and severe dieback and mortality in red and jack pines in Upper Michigan, 402; slash and loblolly infected with *Cronartium fusiforme* galls serve as inoculum for infection of young water oak leaves by aeciospores in Missisippi, 388
- banksiana: new host for brown spot needle blight caused by *Lecanosticta acicola*, and first report of this fungus in Canada, 208
 - contorta: comandra rust, evaluation of antibiotics for control, 528
 - --- var. latifolia: 208; white spored *Peridermium stalactiforme* in Alberta, Canada, 144
 - elliottii var. elliottii: development of pycnia and aecia in albino form of *Cronartium fusiforme* on inoculated seedlings, 842; incidence of *Polyporus tomentosus* in plantations in southeastern United States, 527
 - lambertiana: root disease control by soil fumigation in forest nurseries of California, 245; white pine blister rust, evaluation of basal stem application of cycloheximide and phytoactin for control in Oregon and California, 554
 - patula: 181
 - ponderosa: experimental inoculation with western gall rust for resistance studies, 550; root disease survey to determine extent of damage in natural forest stand shows that *Fomes annosus* is greater menace in California than past reports indicate, 439
 - --- var. scopulorum: root-knot nematode possibly associated with decline in New Mexico, believed to be first report of *Meloidogyne* (may be undescribed species) on this host, 126
 - radiata: *Dothistroma pini* on, first report of this fungus in Chile, 280
 - resinosa and *P. rigida*: *Fomes annosus* infection of stumps effectively prevented by creosote, urea, and liquid borate after 10 months, 22
 - strobus: blister rust cankers not controlled in eastern white pine by cycloheximide and phytoactin antibiotics as reported in western white pine, 224
 - taeda: *Phytophthora cinnamomi* and *Pythium* spp. associated with decline, 1st report of *P. cinnamomi* associated with pine in Louisiana, 596; stumps inoculated with *Fomes annosus* resulted in increased penetration under induced drought conditions, 747
- Pistacia* spp.: new hosts for *Xiphinema* index, 27
- Plasmodiophora brassicae*: 133
- Pleurotus ostreatus*: 552
- Plum (see also *Prunus*): line pattern virus symptoms on indicator trees from virus-infected plum trees, 278
- Podosphaera leucotricha*: 645
- Polyporus lucidus*: 945; (*Ganoderma lucidum*), associated with death of honeylocust in Mississippi, 243
- tomentosus: 360; 527
- Pome Fruit Virus Subproject (IR-2): an interim report, 461
- Potato: black-leg and tuber rots, investigations in Arizona revealed inter-

- action between *Erwinia* sp. and *Pythium aphanidermatum*, 414; Irish, 681; seed production experiments in the United Arab Republic, 761
- Powdery mildew: of apple and bean, controlled by 2 new sulfites, 645; combined control of mildew and blight of carrots, 594; on mock orange, reported for first time from Georgia and the Southeast, 306; new records of *Erysiphaceae*, 709; of papaya, susceptibility of senescent leaves of fruiting plants, 519; of snap bean, several fungicides tested for control, captan + tribasic copper sulfate gave excellent results, 847
- Pratylenchus penetrans*: 923
- *vulnus*: 858
- Prune (see also *Prunus*): bioassay of Italian variety by branch-injection indicate that 8-quinolinol phosphate and 8-quinolinol acetate warrant further investigation as systemic fungicides for orchard trees, 773
- Prunus* spp. (see also common names): ringspot virus complex from plum causes atypical reaction on cherry indicator tree, 278; seven viruses tested for transmission through *Prunus* seed, 902
- Pseudomonas glycinea*: 560; identification of 7 pathogenic races, 557
- *maculicola*: 192
- *phaseolicola*: 29; 110; 238; 371
- *solanacearum*: 275; 520; 836
- Pseudoperonospora cubensis*: 340
- Pseudopeziza ribis*: 196
- Pseudopythophthorus* spp.: 45
- Pseudotsuga menziesii*: *Rhabdocline* needle cast, needle weight determination method accurately estimates intensity of disease, 406; *Rosellinia* needle blight on seedlings, first report of *Rosellinia herpotrichioides* in California, 249
- Psylla pyricola*: 270
- Puccinia coronata* var. *avenae*: pathogenicity of isolates collected in USA, 1961-1965, 935
- *graminis*: method of separation and identification of isolates from mixtures on differential host varieties, 312; physiologic races in the United States in 1964, 332
- --- *f. sp. avenae*: 571; 631
- --- var. *phlei-pratensis*: 849
- --- *secalis*: storage of pycniospores in liquid nitrogen, 576
- --- *tritici*: 299; 673; storage of urediospores in liquid nitrogen, 502
- *recondita*: 108
- --- *f. sp. tritici*: 7th revision of International Register of Physiologic Races of *Puccinia recondita f. sp. tritici*, 756
- *sorghii*: 119
- *striiformis*: 205; 806
- *zoysiae*: 839; 885; found in Mississippi, 770
- Puerto Rico: 484
- Purple stain: caused by *Cercospora kikuchii*, relation of soybean seed infection to duration of pre-flowering, flowering, post-flowering, and pod-maturity periods, 464
- Pyrenochaeta terrestris*: 469
- Pythium* spp.: 393; 596; 695; detection of in soil, 791
- *aphanidermatum*: 414; demonstrated to be pathogenic on barley, oats, rye, wheat and sorghum for the first time, 292
- *debaryanum*: 614
- *ultimum*: 614
- Quarantine: interception in 1965-66 season by plant quarantine inspectors of 3 tree peony species from Japan and of *Lilium regale* bulbs from the Netherlands infected by the nematode *Aphelenchoides fragariae*, 798; proposed isolated laboratory for study of plant pathogens, especially viruses and their vectors, 803
- Quince: virus and virus-like diseases in California, 211
- Races: new physiologic race of *Bremia lactucae* seriously affects 3 previously immune lettuce varieties in south Texas, 460; five para rubber tree clones designated as differentials for identifying races of *Dothidella ulei*, 187; Race 2 of *Fusarium oxysporum f. lycopersici* pathogenic to tomato stocks containing Acc. 160 factor for resistance, 707; serological procedures simplify differentiation of race 1 from race 2 of *Fusarium oxysporum f. sp. lycopersici*, 444; race 1 of *Phytophthora megasperma* var. *sojae* caused rot of soybeans on the High Plains of Texas in 1965, 104; of *Pseudomonas glycinea* on soybean; first report of 7 separate pathogenic races, distinguished by reactions of 7 differential varieties in Minnesota, 557; *Pseudomonas solanacearum* races 1, 2 and 3, serological relationships among, 275; of *Puccinia sorghii*, morphology and pathogenic

- reactions of 2 clonal lines and their inbred recombinants on a homozygous susceptible corn, 119; of sorghums resistant to sooty stripe disease, studies indicate that frequency of resistance is associated with geographic point of origin, 606; evidence for existence of physiologic races of stripe smut attacking Kentucky bluegrass, 430; of *Xanthomonas malvacearum* from Australia, 323
- : rust: of flax rust in the Canadian prairies and in North Dakota in 1963 and 1964, 62; of oat crown rust, 935; 18 oat crown rust races new to North America isolated during recent surveys in Canada, 750; of oat stem, identification of a new and virulent race, 6AFH, from 4 collections of oat stem rust made in Uniform Oat Rust Nursery, 571; oat stem, in a test of select varieties and lines of oats for resistance to stem rust races, adult plants of CI 3034 was most promising source of resistance to 6AFH in hexaploid oats, 631; of *Puccinia graminis*, in the United States in 1964, 332; race 56 of *Puccinia graminis tritici* constituted more than 97% of the pathogen population in wheat in 1960-62, 90% in 1963, 673; seventh revision of the International Register of Physiologic Races of *Puccinia recondita* f. sp. *tritici*, 756; new stripe rust race pathogenic to 'Suwon 92' wheat, 205
- Radiation: gamma, used on oranges in Israel to prolong storage life of fruit, 874
- : ultra-violet, did not stimulate perithecia formation in otherwise non-perithecial cultures in tests for sexual expression in *Colletotrichum orbiculare*, 871
- Ramulispora sorghi*: 606
- Red thread: found on Bermuda grass, 525
- Registration-certification program: organization, history, and progress of Arizona's cooperative system for development of virus-free citrus budwood, 145
- Reservoir hosts: 5 *Nicotiana tabacum* varieties maintained highly infectious cucumber mosaic virus inoculum after 1 1/2 years, 79
- Resistance: cantaloup introduction, PI 140471, approaches immunity to gummy stem blight in the field, 661
- Rhabdocline pseudotsugae*: 406
- Rhamnus cathartica*: mosaic disease on ornamental plantings found to be caused by cucumber mosaic virus, 613
- Rhizoctonia* spp.: 172; 393; 695
- solani: 301; 340; 435; 484; 578; 614; 622; 852; isolated from cotton seed that were acid delinted and surface sterilized with alcoholic mercuric-chloride, 770; role in obtaining and maintaining an adequate tomato stand for production in south Florida, 124
- leaf blight: of azalea, report on disease and control, 354
- Rhizopus stolonifer*: 565
- Rhopalosiphum maidis*: 862
- padi: 380
- Ribes alpinum*: anthracnose, variation in *Pseudopeziza ribis* isolates from several hosts, 196
- Rice: grassy stunt virus disease, transmission by the planthopper *Nilaparvata lugens*, 453; new host of *Curvularia verruculosa*, 314
- Root: inhabiting fungi associated with birdsfoot trefoil, 753
- diseases: in California forest nurseries, control by soil fumigation, 245; disease survey at Boggs Mountain State Forest shows that *Fomes annosus* is greater menace in California than past reports indicate, 439
- knot: considerably reduced in Indian tomato and okra fields by addition of oil-cake amendments to the soil, 668
- Rose: hybrid tea, spotted anthracnose of petals caused by *Spacheloma rosarum*, 934
- Rosellinia herpotrichioides*: 249
- Rot: *Aspergillus* crown, of peanut, serious outbreak in irrigated fields in New Mexico in 1965, 175
- : bacterial soft rot: of Iceberg (Great Lakes) lettuce in Florida Everglades, 938; of pear fruits and coriander leaves, new hosts for *Erwinia aroideae*, 701
- : black core, of mandarin oranges, caused by *Alternaria tenuis*, 127
- : blue-mold, of apple, fungicides for control, 913
- : brown: of citrus, study of length of incubation period of *Phytophthora citrophthora* under natural conditions, 251; 3 *Phytophthora* spp. associated with brown rot syndrome in Western Australian citrus orchards, 494
- : brown stem, of soybeans in Mexico, 598
- : collar, of 1 to 7 year old apple trees propagated on East Malling II and

- Malling Merton 104 rootstocks, 267
- : cotton boll, method for estimating losses, 254
- : foot, of winter wheat, fall applications of diuron to control weeds reduced incidence and severity of, 852
- : fruit, of apple, caused by *Phialophora malorum*, 1st report in Pennsylvania, 701
- : heart, comparison of 2 methods for experimentally inoculating Nuttall oaks, 552
- : leather, of strawberry, occurrence despite lack of rainfall attributed to heavy dew and fog conditions in central California coast, 283
- : perennial canker, of apples in West Virginia and Pennsylvania, 348
- : *Phytophthora*, of soybeans, on the High Plains of Texas in 1965, 104
- : *Rhizopus*, of peaches, preharvest spray experiments in Colorado, 565
- : root: of alfalfa in Ontario, fungi associated with, 695; *Phytophthora* root rot, of alfalfa, found in Canada for first time, 451; identity and relative importance of major root rot pathogens of snap beans in Wisconsin, 578; 1965 survey in Georgia to determine prevalence of organisms in ornamental nurseries and to determine which fungi were involved, 393; *Fusarium*, easy method for evaluating hundreds of lines and varieties of wheat at maturity for resistance, 26; of pea, study of chemical control using methionine and Dexon, divergence between results from field and greenhouse tests, 91; of pine, survey of distribution and severity in Illinois, 446; of red pine seedlings, study of nematode-fungi interaction in soil, 545; caused by *Polyporus lucidus*, tests of pathogenicity on important shade and timber trees of Mississippi Delta, 945; of *Taxus*, 13 diverse *Phytophthora cinnamomi* isolates varied in degree of pathogenicity to 6 cultivars, 84; of tomato, 124
- : ---: *annosus*, significantly higher incidence of infection of *Pinus* spp. revealed by core sampling than by field diagnosis, 589
- : root and butt: distribution of *Fomes annosus* in natural forests of California, 832; red root and butt, incidence of *Polyporus tomentosus* fruiting on infected slash pine trees in a Georgia plantation, 360
- : soil, of cucumber, control, 340
- : stalk: of corn, reaction of inbred lines, varieties, and hybrids to 4 causal fungi, 401; of sorghum, caused by *Aspergillus niger*, intensified by addition of *Macrophomina phaseoli*, 352
- : stem, of geranium, satisfactory technique for checking presence of bacteria in commercial plants, 515
- : stem-end, of Florida lemons for fresh fruit market, 860
- : *Thielaviopsis basicola*, of gloxinia rhizomes from Europe, 709
- Rotenoid: concentration in *Tephrosia vogelii* not affected by fungicides, 484
- Rotylenchulus reniformis*: 149; 500
- Rubber plant: "leaf stripe" disease, foliar nematode shown to be causative agent, 689
- Rubbertree: (para), 5 clones designated as differentials for identifying races of *Dothidella ulei*, 187
- Russetting: of McIntosh and Cortland apples, possibly physiological disease, 151
- Rust: first report of *Uredo ignava* in the United States, found on bamboo in Florida, 112; morphological evaluation of 2 pathogenic clonal lines of *Puccinia sorghi* and their inbred recombinants on a homozygous susceptible corn, 119; seventh revision of the International Register of Physiologic Races of *Puccinia recondita* f. sp. *tritici*, 756
- : blister, of eastern white pine, not controlled in the Lake States by cycloheximide and phytoactin, contrary to results on western white pine, 224
- : comandra, on lodgepole pine, evaluation of antibiotics for control, 528
- : crown, of oat, pathogenicity of isolates collected in USA, 1961-1965, 935; 18 oat crown rust races new to North America isolated during recent surveys in Canada, 750
- : flax, 16 races in varying frequencies in Canadian prairies and in North Dakota, 1963 and 1964, 62; serious reduction of crop value in Texas, 840
- : fusiform: development of pycnia and aecia in albino form on pine, 842; sporophores of *Polyporus tomentosus* var. *circinatus* associated with basal cankers caused by fusiform rust fungus, 527; study of conditions essential to teliospore germination and sporidial production in South Carolina, 432

- : leaf: effect of inhibiting growth of 14 wheat varieties with Cyco-cel on leaf rust infection types, 108; of timothy, 849
- : --- and stem, of timothy, in field and greenhouse screening of 100 Phleum introductions, selections out of PI's 251846, Italy and 267063, Poland were completely resistant -- good resistance found in selections out of 16 other accessions, 849
- : stem: of oats, a new and virulent race, 6AFH, important because virulence for all described genes has been combined into a single phenotype, 571; of oats, varietal reaction to races 6AF and 6AFH reveals CI 3034 as new source of resistance to 6AFH in adult plants in Minnesota, 631; physiologic races of *Puccinia graminis* in the United States in 1964, 332; of timothy, 849; of wheat, inhibition by new antifungal antibiotic, 299
- : stripe: of grasses, systemic activity of 2 new fungicides, 624; new race pathogenic to 'Suwon 92' wheat, 205; of wheat, effective chemical seed treatment for systemic control of seedling infection, 806
- : sweetfern blister: amendments to records of distribution and hosts, 744; occurrence in Alberta and Northwest Territories could be serious threat to hard pine nurseries and plantations, 419
- : western gall, antibiotic tests for control on lodgepole and ponderosa pines, 741; experimental inoculation of ponderosa pine for resistance studies, 550
- : wheat, rapid method of assaying pathogenic potential of populations of *Puccinia graminis tritici*, 673
- : wheat stem, evaluation of nickel + dithiocarbamates for control, 373
- : wheat stripe, mist culture apparatus used for root development studies, 369
- : white pine blister, of sugar pine in Oregon and California, evaluation of basal stem application of cycloheximide and phytoactin for control, 554
- : white spored *Peridermium stalactiforme*, on lodgepole pine in Alberta, Canada, 144
- : zoysia: found in Alabama for first time, 885; in Georgia, 839
- Rye: pathogenicity of *Pythium aphanidermatum*, 292
- Sambucus chinensis*: new host for *Curvularia geniculata*, 488
- Sanitation: 10 recommendations for control of angular leaf spot of cotton that successfully eradicated *Xanthomonas malvacearum* from California, 168
- Scab: pecan, four new fungicide formulations tested for control, 599
- Scleroderris lagerbergii*: 402
- Sclerospora sorghi*: 641
- Sclerotinia borealis*: 69
- *camelliae*: 525
- *homeocarpa*: 770
- Sclerotium bataticola*: 401
- *rolfsii*: 443; 855
- Seed: potato, production experiments in U.A.R., 761; *Rhizoctonia solani* isolated from cotton seed, 770; size of Funza barley seed has significant effect on smut percentage, 101
- : transmission: of latent viruses in commercial peach seed, 740; of psorosis virus through citrus hybrid at levels of 15 to 31%, 81; western X-disease of chokecherry not definitely found to be carried in seed, however nursery seed should come from healthy trees since only low percentage from infected trees will germinate, 659
- treatments: effective chemical treatment for systemic control of seedling infection of wheat by stripe rust, 806; effect on soil-borne fungi at the hypocotyl and cotyledonary surfaces of peanut seed, 162; fungicide evaluations for control of damping-off of table beet seedlings indicate that if PCNB is combined with a standard material used for control of *Pythium* diseases, *Rhizoctonia* in the soil, which varies greatly from year to year will also be controlled, 614
- Septoria linicola*: 840
- Serodiagnosis: use of disposable leaf presses for obtaining plant sap for, 822
- Serological studies: relationships among races 1, 2 and 3 of *Pseudomonas solanacearum* and 4 other plant pathogenic bacterial species, 275
- Serum: anti-, against *Corynebacterium michiganense* can aid in detection of tomato bacterial canker, 96; anti-, against *Pseudomonas solanacearum* races 1, 2 and 3 and a study of their relationships, 275; anti-, against several isolates of *Pseudomonas solanacearum* and the serological

- detection of the fungus in peanut plants by a rapid agar-diffusion method which distinguishes bacterial wilt from wilts caused by other organisms, 836
- Sexual expression: in *Colletotrichum orbiculare*, 871
- Slugs: transmission of cabbage leaf spot caused by *Alternaria brassicicola* by, in France, 764
- Smut: increase in susceptibility of corn inbred W22 by adding potassium to soil associated with decrease in reducing sugar and increase in nitrogen in the pith tissue, 12; survival of various smut fungi cultures for 11 years under mineral oil, 800
- : flag and stripe, systemic activity of 2 new fungicides, 624
- : head, of corn in Idaho, 215
- : stripe: of Kentucky bluegrass, evidence for existence of physiologic races, 430; prevalence, pathogenicity and control studies in New Jersey turf and forage grasses, 294
- Snow mold: on 4 bluegrass varieties, 3 chemicals tested for control during 3 winter seasons in Alaska, 69; further evidence that *Fusarium nivale* is a primary root-rot pathogen of cereals and that in sequence with *Bipolaris sorokiniana* it causes damage in warm as well as in cold seasons in winter wheat in Washington, 154
- Soil: detection of *Pythium* in, 791; induced moisture soil effect on growth of *Fomes annosus* in inoculated loblolly pine stumps, 747; isolating *Verticillium* from, 897; textural studies of in relation to development of hazard ratings for *Fomes annosus* in pine plantations in Virginia, 510
- : amendments, for the control of onion infection and bulb rotting, 469
- : fumigants, for control of *Verticillium* wilt of peppermint, most effective treatments controlled for 3 consecutive crops, 97
- : treatments: combinations tested for control of 2 *Heterodera* spp. occurring with club root organism in Brussels sprouts, 133; effects of chemicals and mulching on development of new apple trees when used at planting time, 201; root disease control by fumigation in California forest nurseries, 245
- Sooty stripe disease: testing for resistant races of sorghums in Nigeria, 606
- Soreshin: 124
- Sorghum: *Colletotrichum graminicolum* spreads through dead stalk tissue cells and is inhibited by living cells, 287
- : downy mildew, part of world collection and Nigerian collection of grain sorghums screened for resistance in 1964 and 1965, 641
- : ergot, host range and epidemiology in Nigeria, 828
- : *Pythium aphanidermatum*, pathogenicity of, 292
- : sooty stripe, studies in Nigeria of the world collection of sorghum indicate that frequency of resistance is associated with geographic point of origin, 606
- : stalk rot, caused by *Aspergillus niger*, intensified by addition of *Macrophoma phaseoli*, 352
- South Carolina: 432; 482
- South Dakota: 819; 917
- Soybean: bacterial blight, inoculation procedures for comparing reaction of soybeans to pathogen, 560
- : brown stem rot, in Mexico, 598
- : cyst nematode, comparative rates of reproduction on 12 hosts, 608
- : *Phytophthora* rot, found on the High Plains of Texas in 1965, 104
- : *Pseudomonas glycinea*, 7 pathogenic races identified in Minnesota, 557
- : purple stain in seed coats found to be closely related to length of flowering stage in selected commercial varieties and plant introductions, 464
- : tobacco ringspot virus of, losses caused by, 910
- Sphacelia sorghi: 828
- Sphaceloma rosarum: 934
- Sphacelotheca spp.: 800
- reiliana: 215
- Spore trap: simple, inexpensive, continuous sampling, 906
- Squash (see Cucurbits)
- Stalk rot: of corn, there was little difference between 3 rating methods evaluated and disease means were not much affected by high N fertilization levels, 655
- Stem canker: oak strain of *Fusarium solani* can infect young oak stems and is more virulent than other strains tested, 160
- Stemphylium botryosum f. lycopersici: 635
- Stinkhorn fungus: unusual occurrence on golf turf, 840
- Storage: lenticel injury to apples resulting

- from fruit packed in molded pulp trays containing urea-formaldehyde wet-strength resins, 885; of pycniospores of *Puccinia graminis secalis* in liquid nitrogen, 576; storage period of oranges extended in Israel by gamma radiation which either delayed incubation of *Penicillium* fungi responsible for decay or prevented their development, 874; of urediospores of *Puccinia graminis tritici* in liquid nitrogen, 502
- Strains: evidence that common pea mosaic virus is a strain of bean yellow mosaic virus, 337; of maize dwarf mosaic virus, 2 distinct strains in northeastern United States identified by host reaction and serological relationships, 814; new strain of tobacco mosaic virus in India, 612; of *Rhizoctonia solani* and *Thielaviopsis basicola* which cause cotton seedling diseases, relation between pathogenicity and sensitivity to chemicals, 301
- Strawberry: new disorder of fruit, caused serious losses in Salinas-Watsonville district of California, 327
- : angular leaf spot, found in Wisconsin for first time, 167
- : leaf spot, relative resistance of 63 varieties and selections studied at 2 locations for 2 years, 105
- : leather rot, occurrence during the absence of rainfall attributed to heavy dew and fog conditions combined with cool temperatures in central California coast, 283
- : *Phytophthora fragariae*, race A-5 found in Maryland and New Jersey, this is the first report in commercial strawberry plantings in the eastern United States, 351
- : viruses: rapid spread into new plantings in western Washington, 586; crinkle virus, over 50 chemical compounds tested did not inactivate virus when sprayed on foliage of infected alpine hybrid vesca plants, 843; virus infection indexing, a simple, efficient, and inexpensive method, 426
- Streptomyces anandii*: 299
- Sugarbeet: nematodes, evaluating varieties and breeding lines for tolerance to nematode invasion, no outstanding resistance was observed, 650
- : viruses: cucumber mosaic virus infectivity increased in vitro on infected leaf tissue, 639; savoy and yellow vein viruses in Great Plains, 1963-64-65, root weight and percentage of sucrose significantly reduced, 457
- Sugarcane: leaf scald, found in Puerto Rico, first report of occurrence in one of the larger islands of the Antilles, 569
- Sunflower: Mexican, wilt of, caused by celery-wilt *Fusarium*, 391
- Surfactants: effect on fungicides applied to watermelon foliage, 327
- Sweetgum: 945
- Sweetpotato: internal root necrosis of New Zealand's 'Owairaka Red', 730; varietal reaction of 24 selections to reniform nematode, 500
- Systemma acicola*: 208
- Tanzania: 593
- Tar spot: of citrus, second report in Florida, chemical control, 357
- Taxus: 13 diverse *Phytophthora cinnamomi* isolates varied in degree of pathogenicity to 6 cultivars, 84
- Teliospore: germination of *Cronartium fusiforme*, 432
- Temperature: of citrus groves, influence on efficacy of hot-water treatment for *Phytophthora citrophthora* control of fruit rot, 867; effect of low postinoculation temperature on colonization of lima bean stems by *Phytophthora phaseoli*, 734; a factor in lenticel injury in apples, 885; hot-water treatments effective in eliminating *Pratylenchus vulnus* from grapevine roots, however soil treatments should still be used, 858; *Thielaviopsis basicola* development enhanced by temperature stress on roots of greenhouse-grown birdsfoot trefoil, 825
- Tennessee: 427
- Tephrosia vogelii*: field evaluations at Mayaguez, Puerto Rico of 8 fungicides for effect on stand count, plant height, dry weight of leaves and stems, leaf:stem ratio, and rotenoid concentration in leaves, 484
- Texas: 7; 75; 104; 149; 460; 645; 840
- Thielaviopsis basicola*: 301; 709; growth on birdsfoot trefoil in the greenhouse increased by temperature stress, 825
- Timothy: screening of introductions for sources of resistance to stem and leaf rust among *Phleum* species, 849
- Tobacco: diseases in Kentucky in 1965, 218
- : cyst nematodes, control with volatile and nonvolatile nematocides, 507
- : false broomrape, field observations

- on response of Maryland tobacco to, 627
- : viruses: cucumber mosaic virus, highly infectious inoculum obtained from Kentucky 57, 304-1, T.I. 245, Holmes Samsoun and Xanthi-nc varieties after 1 1/2 years, 79; survey of virus diseases for distribution and relation of previous cropping history to occurrence in North Carolina, 308
- Tomato: 681; unusual occurrences of late blight, buckeye rot, leaf mold and bacterial spot in Louisiana, 686
- : bacterial canker, 3 serological diagnostic methods for, 96
- : bacterial spot, analysis of role in production in south Florida has resulted in effective control program using copper, maneb, and streptomycin, 699
- : curly top virus, wooly and dwarf tolerance to, Colorado, 309
- : foliage diseases, severity affected by mode of irrigation in Israel, 635
- : *Fusarium oxysporum* f. sp. *lycopersici*, serological procedures simplify differentiation of race 1 from race 2, 444
- : nematodes, 'Rutgers', used as indicator in nematocide dose and exposure time tests, 915
- : *Rhizoctonia solani*, role in obtaining and maintaining an adequate tomato stand for production in south Florida, 124
- : southern blight, losses reduced, and fruit yields and plant vigor improved by deep plowing, 443
- : wilt, every major tomato-growing area of Florida resurveyed for race 2 of *Fusarium oxysporum* f. *lycopersici* which is pathogenic to stocks containing Acc. 160 factor for resistance, 707
- Top damping-off: of tomato, 124
- Toxin: production in vitro by non wilt-inducing isolates of *Fusarium*, 261
- Transmission: apple virus transmission studies with selective *Malus* indicators, 40; bud transmission of dwarfing virus in sweet cherry, 367; of cabbage leaf spot by slugs in France, 764; graft, of a ringspot-like virus from white ash to herbaceous plants, 905; of latent viruses in commercial peach seed, 740; of seven viruses through *Prunus* seed tested, 902
- Trichoderma viride*: 747
- Trinidad: 478
- Tung: angular leaf spot: effect on yield and oil content of fruit from trees sprayed with Bordeaux mixture or petroleum oil for 5 years, 54; evaluation of various fungicides for control, 717
- Tylenchulus semipenetrans*: 822; 960
- United Arab Republic: 761
- Uredo ignava*: 112
- Ustilago* spp.: 800
- maydis: 12
- nuda: 101
- striiformis: 294; 430
- Vectors associated with: barley yellow dwarf virus, on wheat -- *Rhopalosiphum padi*, 234
- : corn stunt virus, -- corn leafhopper, gamagrass found to be new host for, 570
- : cucumber mosaic virus, of *Rhamnus cathartica* -- *Myzus persicae*, 613
- : decline virus of pear -- pear psylla, 270
- : grassy stunt disease of rice, -- planthopper, *Nilaparvata lugens*, 453
- : maize dwarf mosaic virus of corn -- corn leaf aphid, 862
- : oak wilt -- small oak bark beetle, 45
- : strawberry viruses -- strawberry aphid, 586
- : streak mosaic, of wheat -- *Aceria tulipae*, 917
- : stunt (new virus disease) of peanut -- green peach aphid, 139
- : watermelon mosaic virus -- melon aphid, 305
- : wheat streak mosaic virus: of corn -- *Aceria tulipae*, 819; of corn and wheat -- wheat curl mites, 951
- Verticillium* spp.: method for isolating *Verticillium* from field soil, 897
- albo-atrum: 930
- dahliae (=V. albo-atrum): 97
- Vigna sesquipedalis*: intercellular colonization and the production of pectic enzymes by Malayan isolates of *Fusarium*, 116
- Vinca minor*: dagger nematodes can cause severe decline under some circumstances, 420
- Virginia: 139; 510
- Virus, Virus disease: availability of "Index of Plant Virus Diseases", 624; establishment of virus-free citrus budwood through Arizona's Cooperative Citrus Registration-Certification Program, 145; possible cause of an internal root necrosis of a New Zealand sweetpotato, 730;

- first report of a virus resembling tobacco streak in western United States isolated from dodder growing on a desert shrub in southern California, shrub may be symptomless reservoir for virus, 479
- : of apple: preliminary report of IR-2 Pome Fruit Virus Subproject stating objectives and procedures necessary for virus indexing of latent diseases, 461; transmission studies with selective *Malus* indicators, 40
- : apricot ring pox, 91 foreign varieties found to be hosts, 289
- : aster yellows, susceptibility of 44 grasses tested, host range now includes 8 grasses (including oats), 17
- : barley yellow dwarf: report of recovery tests on 10 weed species from Mississippi suspected to be natural summer hosts for BYDV, 692; resistance in selected barley varieties, 380; 12 wheat varieties susceptible to, in Kansas, 234
- : of beans: tests of bean cultivars and breeding lines for sources of resistance to bean yellow mosaic, alfalfa mosaic, western ringspot, and common bean mosaic viruses, 532; common pea mosaic strongly indicated as strain of bean yellow mosaic, 337
- : calico, of lettuce, caused by alfalfa mosaic not by tobacco ringspot, 629
- : of *Cattleya*, simple method for detection of a flexuous-rod virus associated with infectious *Cattleya* blossom necrosis, 779
- : of cherry, bud transmission of dwarfing virus in sweet cherry, 367
- : common pea mosaic, strongly indicated as strain of bean yellow mosaic, 337
- : of corn: distribution and identification of mechanically transmissible corn viruses in Louisiana, 325; occurrence of mechanically transmissible virus of corn in Mississippi, 409; evidence that streak disease of corn in California is caused by sugarcane mosaic and that corn stunt does not occur in that State, 318; corn stunt, corn leafhopper found on new host, gamagrass, in Mississippi, 570
- : of cranberry, virus-like ringspot symptoms on fruits and leaves of Searles and Howes varieties in Wisconsin, 543
- : crinkle, of strawberry alpine hybrid vesca, studies on inactivation by chemotherapy, 843
- : cucumber mosaic: causes mosaic disease of canna, 899; causes mosaic disease of *Rhamnus cathartica* in Wisconsin, 613; 5 *Nicotiana tabacum* varieties serving as reservoir hosts maintained highly infectious inoculum after 1 1/2 years, 79; from sugarbeet, in vitro increase in virus infectivity, 639
- : cucurbit viruses, on the west coast of Mexico, 947
- : curly top, tolerance of woolly and dwarf tomatoes to, 309
- : grassy stunt, of rice, transmission by planthoppers, 453
- : green ring mottle virus, of sour cherry, severe fruit necrosis caused by strains of, 59
- : maize dwarf mosaic: a heavy-duty, mobile air blast inoculator has been designed, built, tested, and proved to be rapid and efficient for both field and greenhouse use, 363; 2 distinct strains occurring in northeastern United States, 814; presence established in several counties in Pennsylvania in 1965, 6
- : ---: of grasses: mechanical inoculation of St. Augustine grass seedlings resulted in 80% infection, 441; first report of maize dwarf mosaic on gamagrass, Illinois, 65; positive transmission of one isolate from Johnson grass in Kansas, first report, 1965, 11
- : ---: of corn: found in Iowa for first time, 88; host range, properties, and partial purification of a Missouri isolate, 862; in Tennessee, 427
- : of Malvaceous plants, chemo- and thermotherapy for control of leaf curl, 88
- : mule's ear, of peach, 64
- : muskmelon necrotic fleck, new strain of tobacco mosaic virus isolated from muskmelon in Arizona, 49
- : of papaya, relationship of disease with distance and time within a field, 908
- : of peach, transmission of latent viruses in commercial peach seed, 740
- : of peanuts: new disease (stunt) causes severe dwarfing and fruit suppression, mechanical, graft, dodder, and aphid transmission, 139; severe stunting and yield reduction, and obvious spread of unidentified virus in 5 North Carolina locations, 136

- : pear decline, new brown leaf-vein symptom associated with infected greenhouse-grown pear plants, 270
- : of pepper, oil sprays delay field spread but do not increase yields, 158
- : phloem necrosis, (*Morsus ulmi*) of American elm in Mississippi Delta, 751
- : potato, seed production experiments in U. A. R., 761
- : of Prunus: 7 viruses tested for transmission through Prunus seed, 902; Prunus ringspot complex from plum causes atypical symptoms on Shiro-fugen flowering cherry, 278
- : psorosis virus, transmitted through seeds of at least one citrus hybrid ('Carrizo' citrange) at levels of 15 to 31%, 81
- : of quince, in California, 211
- : sour cherry yellows, of cherry, effect of gibberellic acid sprays on fruit spur development of infected trees at different growth stages and environmental conditions, 240
- : of strawberry: rapid spread into new plantings in western Washington, 586; a simple, efficient, and inexpensive method of collecting leaves for virus infection indexing, 426
- : of sugarbeet, savoy and yellow vein diseases significantly reduced root weight and sucrose percentage in Great Plains, 1963-65, 457
- : sugarcane mosaic, reaction of 180 sweet and dent corn varieties to Johnson grass strain of, 793
- : of tobacco, viruses present, relation of occurrence to previous cropping history and distribution in North Carolina, 308
- : tobacco mosaic, new strain naturally attacks chilli pepper variety Puri orange in India, and has a wide host range, 619
- : tobacco ring spot: of guar, new occurrences in Texas and Oklahoma, 7; of soybean, losses caused by, 910
- : watermelon mosaic, aluminum foil for repellency to melon aphids, 305
- : western X-disease, chokecherry infected in central and northern Great Plains, study of possible seed transmission of the disease, 659
- : wheat streak mosaic: epidemiology in South Dakota, 1962-66, and spread of virus under field and greenhouse conditions, 917; tests indicate that Ohio 3A virus culture is strain of wheat streak mosaic and its occurrence in Ohio suggests that wheat and susceptible corn could be year-round hosts throughout corn belt, 951; of corn, host range and spread of virus and vector in South Dakota, 1962-66, 819
- : of white ash, graft transmission of a ringspot-like virus from white ash to herbaceous plants, 905
- Wart disease, of subterranean clover caused by *Physoderma trifolii*, first report on this host in the United States, 797
- Washington: 154; 586; 885
- Watermelon (see Cucurbits)
- Weather: freeze of November 1955 in Pacific Northwest may be predisposing factor in severe dieback of *Ceanothus* spp., 399; frost injury on wheat, 928; temperature-relative humidity-time technique for forecasting *Cercospora* leafspot of peanut, 810
- West Virginia: 348; 545
- Wheat: frost injury, 928; pathogenicity of *Pythium aphanidermatum*, 292
- : *Cephalosporium* stripe disease, first published report from Michigan, 190
- : rots: common root rot, easy method for evaluating hundreds of lines and varieties at maturity for resistance, 26; fall applications of diuron to control weeds reduced incidence and severity of foot rot of winter wheat caused by *Cercospora herpotrichoides* and *Rhizoctonia solani*, 852; further evidence that *Fusarium nivale* is a primary root-rot pathogen of cereals and that in sequence with *Bipolaris sorokiniana* it causes damage in warm as well as in cold seasons in winter wheat in Washington, 154
- : rusts: rapid method of assaying pathogenic potential of *Puccinia graminis tritici* populations for use in epidemic prediction and in detection of new, dangerous virulences, 673; seventh revision of the International Register of Physiologic Races of *Puccinia recondita* f. sp. *tritici*, 756
- : ---: leaf, effect of host growth inhibition by Cycocel on infection types, 108
- : ---: stem: evaluation of nickel + dithiocarbamates for control, 373; physiologic races of *Puccinia graminis* in the United States in 1964,

- 332; pustules markedly reduced by new antibiotic, Pentaene G8, 299
- : ---: stripe: effective chemical seed treatment for systemic control of seedling infection, 806; mist culture apparatus used for root development studies, 369; new stripe rust race pathogenic to 'Suwon 92', 205
- : viruses: barley yellow dwarf virus, field tests indicated susceptibility of 12 varieties of winter wheat in Kansas, 234; wheat streak mosaic virus in the Ohio Valley, 951; streak mosaic, epidemiology of virus in South Dakota, 1962-66, and spread of virus under field and greenhouse conditions, 917
- Wilt: of Mexican sunflower, caused by celery-wilt *Fusarium*, 391; of alfalfa in Ontario, fungi associated with, 695
- : bacterial: of peanut, disease reaction and symptom expression of 17 cultivars, 520; *Pseudomonas solanacearum*, detected serologically in peanut plants by an agar-diffusion method, 836; races 1, 2, and 3 detected serologically and compared, 275
- : *Fusarium*: of chrysanthemum, Yellow Delaware cultivar under high N 3 weeks before inoculation more susceptible to wilt than plants under low N, 882; resurvey of every major tomato-growing area of Florida for, 707
- : oak: distribution extended in Arkansas, 862; evaluation of small oak bark beetle as a vector, 45; field evaluation of antibiotics and chemicals for control in northern pin oaks, 736
- : *Verticillium*: comparison of virulence of isolates of *V. albo-atrum* from several of the cotton-growing regions of the U.S., Mexico, and Peru, 930; of peppermint, control by soil fumigation for 3 consecutive crops, 97
- Wisconsin: 167; 192; 539; 543; 578; 677; 957
- Xanthomonas albilineans*: 569
- alfalfae: 731
- fragariae: 167
- malvacearum: 323; eradicated from California through sanitation practices, 168
- pelargonii: 515
- phaseoli: 29; 110; 371
- pruni: 112
- vesicatoria: 635; 686; 699
- Xiphinema americanum*: 420
- index: 27

CROPS RESEARCH DIVISION

Prepared by Sheila D. Thompson

For sale by the Superintendent of Documents, U. S. Government Printing Office, Washington, D. C. 20402 - Price \$4.00 per year, domestic; \$5.00 per year, foreign. Single copy price will vary.

UNITED STATES DEPARTMENT OF AGRICULTURE
Agricultural Research Service
Beltsville, Maryland 20705

Postage and Fees Paid
U.S. Department of Agriculture

Official Business